


# Association between social activity and development of dementia in hearing impairment: A cohort study in Japan from Japan Gerontological Evaluation Study

メタデータ	言語: en 出版者: SAGE Publications 公開日: 2024-03-11 キーワード (Ja): キーワード (En): 作成者: 小島, 香 メールアドレス: 所属:
URL	<a href="http://hdl.handle.net/10271/0002000110">http://hdl.handle.net/10271/0002000110</a>

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.



# Association between Social Activity and Development of Dementia in Hearing Impairment: A Cohort Study in Japan from Japan Gerontological Evaluation Study

Gerontology & Geriatric Medicine  
Volume 8: 1–11  
© The Author(s) 2022  
Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
DOI: 10.1177/23337214221100621  
[journals.sagepub.com/home/ggm](https://journals.sagepub.com/home/ggm)  


Kaori Kojima, MS<sup>1</sup> , Eisaku Okada, PhD<sup>1,2</sup>, Toshiyuki Ojima, MD, PhD<sup>1</sup>, Jun Aida, DDS, PhD<sup>3</sup>, Yoshimune Hiratsuka, MD, MPH, PhD<sup>4</sup>, and Katsunori Kondo, MD, PhD<sup>5,6</sup>

## Abstract

We aimed to clarify the association between social activity and the development of dementia in older adults by hearing-impaired (HI) status. We applied a community-based prospective cohort study over 6 years as part of the Japan Gerontological Evaluation Study. The study included 53,549 participants aged 65 years and older who did not require long-term care. A baseline questionnaire survey was conducted; explanatory variables included physical and social activities, and the objective variable was dementia onset assessed by standardized protocol. Cox regression models were used to calculate hazard ratios (HRs) for dementia stratified by HI status. During the follow-up period, 6013 (11.2%) participants developed dementia. Analyses revealed increased dementia risk for participants with HI who participated in the following activities less than once a month: sport groups (HR 2.17, 95% CI 1.53-3.08), hobby groups (HR 1.70, 95% CI 1.34-2.17), going out (HR 2.19, 95% CI 1.51-3.17), and meeting with friends (HR 1.27, 95% CI 1.06-1.53). HI and lack of social activity increase the risk of dementia. The study results indicate that there is an association between low social activity and the development of dementia in people with HI; the strongest associations were found for low participation in sports and hobby groups.

## Keywords

older adults, dementia, hearing impairment, longitudinal study, dementia risks

Manuscript received: March 2, 2022; final revision received: April 21, 2022; accepted: April 26, 2022.

## Introduction

Hearing impairment (HI) is one of the risk factors for dementia (Livingston et al., 2017). HI, even when mild, increases the long-term risk of developing dementia (Daniele et al., 2018). Among the preventable risk factors for dementia, HI has been identified as the most important modifiable risk factor (Livingston et al., 2017).

In addition to dementia, HI can lead to depression (Li et al., 2014), apathy (Sugawara et al., 2011), and limited participation in daily activities due to reduced stimulus input, negative impact on family relationships and social interactions (Servidoni & Conterno, 2018), social isolation due to

<sup>1</sup>Department of Community Health and Preventive Medicine, Hamamatsu University School of Medicine, Hamamatsu City, Japan

<sup>2</sup>Faculty of Social Policy & Administration, Hosei University, Machida, Japan

<sup>3</sup>Department of Oral Health Promotion, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, Bunkyo-ku, Japan

<sup>4</sup>Department of Ophthalmology, Juntendo University Graduate School of Medicine, Bunkyo-ku, Japan

<sup>5</sup>Department of Social Preventive Medical Sciences, Center for Preventive Medical Sciences, Chiba University, Chiba, Japan

<sup>6</sup>Department of Gerontological Evaluation, Center for Gerontology and Social Science, National Center for Geriatrics and Gerontology, Obu City, Japan

## Corresponding Author:

Toshiyuki Ojima, Department of Community Health and Preventive Medicine, Hamamatsu University School of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu city 431-3192, Japan.  
Email: [ojima@hama-med.ac.jp](mailto:ojima@hama-med.ac.jp)



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE

and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

impaired communication (Strawbridge et al., 2000), and adverse effects on the degree of social interactions (Ogawa et al., 2019). The possibility of a causal relationship between the inhibition of social interaction, depletion of cognitive reserve, social isolation, environmental changes, and a combination of these factors with dementia has also been reported (Ives et al., 1995).

Recently, the importance of social participation in the cognitive functioning of older adults has been widely highlighted. Social participation has been defined by various other researchers (Putnam, 1995; Berkman, 1995). Previous studies have shown that social activity among older adults is associated with reduced risk for long-term care and death (Kanamori et al., 2014; Murata et al., 2017; Saito et al., 2019; Aida et al., 2011; Kawachi et al., 1996).

Older adults with high social participation have higher cognitive function than those with low social participation (Barnes et al., 2004; Holtzman et al., 2004), and social participation suppresses the decline of cognitive function (Uchino, 2006). The strength of social networks, such as social activities and socializing with friends and family, is associated with cognitive functioning and prevents dementia (Holtzman et al., 2004). Decreased social participation increases the risk of developing dementia (Saczynski et al., 2006), and people with fewer social networks have a higher risk of cognitive impairment (Nemoto et al., 2017). Moreover, living alone and having no personal connections doubles the risk of developing dementia in 3 years (Fratiglioni et al., 2000). However, to our knowledge, no study has revealed the relationship between the type of social participation and dementia among older people with HI. In this study, we used large-scale longitudinal data over 6 years to clarify the association between activities performed by older adults with HI and the risk of developing dementia in Japan.

## Methods

### Study Design and Participants

The study design was a community based prospective cohort study from the Japan Gerontological Evaluation Study (JAGES). The baseline survey was conducted from August 2010 to January 2012 by distributing questionnaires to 95,827 randomly-selected older adults who had not been certified as requiring support or long-term care and resided in 13 municipalities in seven prefectures in Japan.

We excluded 7889 people who had missing sex or age information on the self-administered questionnaire, whose answers were inconsistent with the list received from the municipalities, who returned the personal identification code detached from the questionnaire, and who did not give consent for the use of their data for research. Therefore, valid data at baseline was 54,537. We further excluded 153 people with reduced activities of daily living (ADL) who reported difficulty in caring for themselves. We followed up with these people using the “data on the certification of the need for

long-term care” that included the onset of dementia and the “data on the levy of long-term care insurance” that included mortality for 6 years to determine the presence of dementia or the need for long-term care.

To reduce the effect of the reversal of causality, we excluded 835 individuals whose follow-up period was less than 365 days. Deaths within 365 days were also excluded. Finally, 53,549 participants (24,534 men and 29,015 women) were included in the analysis. The follow-up status of the participants is shown in Figure 1.


### Variables

**Outcome Variable.** HI was defined as those who selected self-reported HI as a disease under treatment during the baseline survey.

The outcome variable of the development of dementia was determined if a participant was certified as requiring support or care and classified as IIa (Hisano, 2009; Noda et al., 2018) or higher for “level of independence in daily living for the elderly with dementia” defined by the Ministry of Health, Labor and Welfare. The IIa level is assigned when symptoms, behaviors, or difficulties in communication, related to dementia, that interfere with daily life are observed; the person can be independent if someone pays attention to them. Symptoms of dementia include frequently getting lost, noticeable errors in shopping, office work, money management, and other things that were previously possible. This dementia index is strongly correlated with the Mini-Mental State Examination ( $r = -0.74$ ) (Hisano, 2009). The dementia scale was an effective indicator for the clinical diagnosis; reported sensitivity and specificity were 73% and 96% for rank II dementia, respectively (Noda et al., 2018).

### Explanatory and Covariates Variables

Explanatory variables about social activities included the frequency of activities and the presence of leisure activities. The frequency of going out (twice a week or more, once a week to once a month or more, once a month or less), frequency of meeting friends, and participation in hobby and sport groups (once a month or more) were surveyed (Willie-Tyndale et al., 2016). Covariates to be adjusted included age, sex, self-rated health, ADL, depressive symptoms assessed by geriatric depression screening scale-15 (GDS) (Brink et al., 1982), smoking status, education, equivalent income, marital status, and living arrangement. Age was categorized into five groups: 65–69, 70–74, 75–79, 80–84, and 85 years and older. Self-rated health was categorized into good (very good, fair) and poor (not so good, poor). ADL responses were requested in the following categories: no assistance, partial assistance, and full assistance. Level of depressive symptoms was categorized using the GDS: 0–4 points as no depression, 5–9 points as mild depression, and 10–15 points as depression. Smoking status was categorized into non-smoking (never smoked, quit more than 5 years ago, and do not smoke now)



**Figure 1.** Flow of participants for analytic sample.

and smoking (quit less than 5 years and do not smoke now, still smoking). Equivalent income was categorized into less than 2 million yen, 2–4 million yen, and 4 million yen or over). Equivalent income was calculated by dividing the total annual income of the entire household by the square root of the number of household members. Education was categorized by years of education: less than 10 years, 10 years or more.

### Statistical Analyses

We described the distribution of each variable used in the analysis; the proportion of people who developed dementia by each attribute is expressed in the HI group and normal hearing group, and the chi-square test was used to compare the percentages by each attribute.

To examine the association between social activities and the development of dementia, hazard ratios (HRs) and 95% confidence intervals (95% CIs) were calculated using the Cox proportional hazards model after stratification into two groups: HI and normal hearing. The group with higher participation in each social activity was set as the reference in all the analyses. Samples with missing values in the covariates were included in the analysis by setting the missing categories, i.e. missing-indicator method.

Factors that were significant at the  $p < 0.10$  level in univariate regression were used. In addition, we used factors whose explanatory variables have been suggested to be associated with hearing impairment and dementia in previous studies (Emmett & Francis, 2015; Kochkin, 2007; Yamada et al., 2011; Wayne & Johnsrude, 2015; Uchida et al., 2019;

Panza et al., 2015; Amieva et al., 2018). Covariates of age and sex were included in Model 1, self-rated health, ADL, current smoking, and depressive symptoms, living arrangement, equivalent income, education, and marital status were included in Model 2. We performed analyses using IBM SPSS Statistics version 24. The statistical significance level was set at  $p < 5\%$ .

### Ethical considerations

The JAGES protocol was reviewed and approved by the Ethics Committee on Research of Human Subjects at Nihon Fukushi University (approval No. 1005) and the Ethics Committee of the Chiba University Faculty of Medicine (approval No. 2493).

### Results

The mean age (standard deviation, SD) of the participants was 73.62 (SD 5.78) years. The mean age of participants classified by gender was 73.27 (SD 5.60) years for males and 73.89 (SD 5.89) years for females. The mean follow-up period was 2063 days (2025 days for men and 2095 days for women). The number of participants who developed dementia during the follow-up period was 6013 (11.2%) (2577 [10.0%] men and 3436 [11.8%] women). The demographics of the study subjects were divided according to the hearing status (Table 1). Among the participants, 3369 were HI, and 44,487 had normal hearing. Those who developed dementia were 860 (21.3%) in HI, and 5153 (10.4%) in normal hearing. Significant differences were found for all variables in normal hearing, and almost all variables except for living arrangement in HI. Not shown in the table, the hazard ratio for dementia comparing normal hearing with hearing loss was (1.28, 95%CI 1.19-1.38), adjusted for sex and age. Hazard ratios of the development of dementia by social activities are shown in Table 2. In Model 2 in the HI group, significantly higher HRs were observed for going out less than once a month (2.19, 95% CI 1.51-3.17), participation in sport groups less than once a month (2.17, 95% CI 1.53-3.08), hobby groups less than once a month (1.70, 95% CI 1.34-2.17), leisure activities (1.57, 95% CI 1.31-1.88), and meeting with friends less than once a month (1.27, 95% CI 1.06-1.53). The HRs were higher for sport groups and hobby groups in HI compared with the normal hearing group, while it was almost the same for leisure activities.

### Discussion

In this cohort study, we revealed an association between social activities and the development of dementia in people with HI. This is the first study to show the association in older people with HI to the best of our knowledge. After adjusting for factors such as age, sex, physical factors, mental factors, educational background, and living

conditions, there was an association between social interaction and social activities with the development of dementia in people with HI. Stronger associations were found for participation in sport and hobby groups, which involve deeper interaction and communication.

The crude incidence of dementia was more than twice as high. After adjusting for gender and age, the incidence was 1.28 times higher than in healthy subjects. The results were the same as in the previous study (Michalowsky et al., 2019). Preventing cognitive decline is very important for older people with HI.

Hazard ratio of people with HI were higher for those who participated in sports (HR2.17 vs. 1.28) and hobby (HR1.70 vs. 1.34) groups less than once a month compared with people with normal hearing. For other social activities, the difference between people with normal hearing and HI people was minimal. Leisure activities had a lower HR than hobby and sports groups. This is because leisure activities include pachinko, gardening, painting, and other activities that do not require communication and can be enjoyed individually. Dementia was not associated with participation in social activities once a month. Instrumental Activities of Daily Living (IADL) limitations were reported to be associated with a long-term risk of developing dementia (Bennett et al., 2006). A decline in ADL may be associated with dementia in older adults with HI. For the HI in Table 1, there was no association with living arrangement. Even if HI people live alone, having social activity opportunities and interaction may prevent dementia. Participation in socially interactive activities with others and physical activities is important in reducing the risk of dementia.

Hazard ratio of meeting with friends less than once a month was high in this study. HI, cognitive function, isolation, and depression are interrelated (Dawes et al., 2015). Therefore, social activities may prevent isolation and depression and prevent cognitive impairment for people with HI. Previous studies have reported that social networks prevent dementia in general older people (Levasseur et al., 2010), which is consistent with reports from longitudinal studies, which found that lack of social connections was associated with an increased risk of dementia (Uchino, 2006). Social networks may facilitate access to healthy behaviors for HI people, which in turn may help prevent dementia. In a study that analyzed the relationship between social relationships between HI people and older adults, the researchers found that HI people tended to have smaller networks, and HI was one of the factors associated with social relationships in older adults (Pérès et al., 2008). It is important for HI people to widen their social networks.

Social activities in congenital HI and profound acquired HI are different. There are many reports on social activities in HI people with congenital HI (Moog et al., 2011; Bat-Chava et al., 2005). The use of cochlear implants at an early age has been reported to improve students' participation in club activities, sports meetings, communication, and social skills.

**Table 1.** Characteristics of the Surveyed Participants.

Variables	Categories	Hearing impairment				Normal hearing			
		Dementia n (%)	No dementia (%)	Total (n = 4039)	p-value	Dementia	No dementia	Total (n = 49510)	p-value
Dementia		860 (21.3)	3179 (78.7)	4039		5149 (10.4)	44361 (89.6)	49,510	
Sex	Male	381 (19.9)	1536 (80.1)	1917	<0.01	2196 (9.7)	20421 (90.3)	22,617	<0.01
	Female	480 (22.6)	1642 (77.4)	2122		2957 (11.0)	23936 (89.0)	26,893	
Age	65-69	31 (6.8)	422 (93.2)	453	<0.01	321 (2.3)	13835 (97.7)	14,156	<0.01
	70-74	65 (7.8)	771 (92.2)	836		801 (5.3)	14396 (94.7)	15,197	
	75-79	188 (16.5)	952 (83.5)	1140		1400 (12.5)	9822 (87.5)	11,222	
	80-84	296 (29.4)	673 (70.6)	953		1516 (24.4)	4685 (75.6)	6201	
	85 and older	296 (45.1)	361 (54.9)	657		1115 (40.8)	1619 (59.2)	2734	
Sport group	>1 month	46 (9.2)	454 (90.8)	500	<0.01	537 (6.0)	8437 (94)	8974	<0.01
	≤1 month	552 (22.5)	1904 (77.5)	2456		3169 (10.6)	26685 (89.4)	29,854	
	Missing	262 (24.2)	75.8	1083		1447 (13.5)	9235 (86.5)	10,682	
Hobby group	>1 month	118 (12.5)	825 (87.5)	94	<0.01	1004 (6.6)	14180 (93.4)	15,184	<0.01
	≤1 month	481 (22.7)	1636 (77.3)	2117		2796 (11.2)	22165 (88.8)	24,961	
	Missing	261 (26.7)	718 (73.3)	979		1353 (14.4)	8012 (85.6)	9365	
Going out	>2 times a week	617 (18.8)	2663 (81.2)	3280	<0.01	3999 (9.2)	39395 (90.8)	43,394	<0.01
	>1 times a month	113 (29)	276 (71)	389		569 (20.6)	2196 (79.4)	2765	
	≤1 time a month	46 (51.1)	44 (48.9)	90		180 (35.5)	327 (64.5)	507	
	Missing	84 (30)	196 (70)	280		405 (14.2)	2439 (85.8)	2844	
Meeting with friends	>1 month	481 (18.8)	2080 (81.2)	2561	<0.01	3182 (9.1)	31951 (90.9)	35,133	<0.01
	≤1 month	250 (24.1)	787 (75.9)	1037		1360 (12.6)	9441 (87.4)	10,801	
	Missing	129 (29.3)	312 (70.7)	441		611 (17.1)	2965 (82.9)	3576	
Leisure activities	Yes	279 (15.9)	1478 (84.1)	1757	<0.01	1900 (7.1)	24853 (92.9)	26,753	<0.01
	No	486 (25.6)	1414 (74.4)	1900		2719 (14.3)	16267 (85.7)	18,986	
	Missing	95 (24.9)	287 (75.1)	382		534 (14.2)	3237 (85.8)	3771	
Self-rated health	Good	428 (18.3)	1915 (81.7)	2343	<0.01	3376 (8.5)	36297 (91.5)	39,673	<0.01
	Poor	406 (25.6)	1181 (74.4)	1587		1563 (18.0)	7132 (82.0)	8695	
	Missing	26 (23.9)	83 (76.1)	109		214 (18.7)	928 (81.3)	1142	
Activities of daily living	Independent	750 (20)	2996 (80)	3746	<0.01	4694 (9.9)	42894 (90.1)	47,588	<0.01

(continued)

Table 1. (continued)

Variables	Categories	Hearing impairment				Normal hearing			
		Dementia n (%)	No dementia (%)	Total (n = 4039)	p-value	Dementia	No dementia	Total (n = 49510)	p-value
Current smoking	Minimum assist	73 (46.8)	83 (53.2)	156	0.04	237 (38.7)	375 (61.3)	612	<0.01
	Missing	37 (27.0)	100 (73.0)	4039		222 (16.9)	1008 (83.1)	1310	
	Yes	624 (20.7)	2386 (79.3)	3010		3722 (10.1)	33218 (89.9)	36,940	
Depressive symptoms	No	104 (20.3)	408 (79.7)	512	<0.01	728 (9.8)	6678 (90.2)	7406	<0.01
	Missing	132 (25.5)	385 (74.5)	517		703 (13.6)	4461 (86.4)	5164	
	No depression	235 (18.7)	1023 (81.3)	1258		1762 (7.7)	21028 (92.3)	22,790	
	Mild depression	532 (21.7)	1921 (78.3)	2453		3072 (12.4)	21693 (87.6)	24,765	
Marital status	Depression	83 (27.6)	218 (72.4)	301	<0.01	273 (16.4)	1390 (83.6)	1663	<0.01
	Missing	10 (37.0)	17 (63.0)	27		46 (15.8)	246 (84.2)	292	
	Married	448 (17.9)	2051 (82.1)	2499		2876 (8.2)	32146 (91.8)	35,002	
Living arrangement	Others	412 (26.8)	1128 (73.2)	1540	0.92	2277 (15.7)	12211 (84.3)	14,488	<0.01
	Living alone	116 (21.4)	425 (78.6)	541		781 (12.9)	5277 (87.1)	6058	
	Living with others	744 (21.3)	2754 (78.7)	3498		4373 (10.1)	39079 (89.9)	43,452	
Education	≤10 years	548 (22.1)	1937 (77.9)	2485	<0.01	2820 (12)	20636 (88.0)	23,456	<0.01
	>10 years	251 (18.0)	1143 (82.0)	1394		2024 (8.2)	22712 (91.8)	24,736	
Equivalent income	Missing	61 (38.1)	99 (61.9)	160	<0.01	308 (23.4)	1010 (76.6)	1318	<0.01
	Low	267 (21.7)	965 (78.3)	1232		1467 (12.4)	10407 (87.6)	11,874	
	Middle	207 (17.7)	961 (82.3)	1168		1464 (8.7)	15405 (91.3)	16,869	
	High	175 (19.1)	741 (80.9)	916		1212 (8.5)	13128 (91.5)	14,340	
	Missing	211 (29.2)	511 (70.8)	723		1009 (15.7)	5418 (84.3)	6427	

The total number of participants is shown as a percentage. Comparisons were made using the chi-square test. For missing values, missing categories were created and included in the analysis.

Many people with congenital HI who do not use cochlear implants have learned sign language and use it to communicate with others. They are often socially active within the community of congenital HI. On the other hand, it is difficult for people with profound acquired HI to learn sign language and join the community of HI. Limited communication at home and with friends has been associated with social introversion and loneliness (Knutson & Lansing, 1990). HI has many negative effects on daily life, including depression (Li et al., 2014) and apathy (Sugawara et al., 2011). Social networks have been associated with factors that affect brain function, such as stress and depression (Dori et al., 1994), and neuroendocrine measures of stress have been associated with dementia (Brice & Strauss, 2016).

Thus, social activities may be an effective way to maintain communication opportunities while covering the HI of older adults in the community. People should start social activities such as sport or hobby clubs early, from the time their hearing function was normal and should continue them even if hearing function declines. A longstanding relationship will be useful for mutual understanding in the club. Moreover, the use of hearing aids and consultations with audiologists will be effective for the HI to more deeply interact and communicate with others. Financial assistance for hearing aids and the development of better hearing aids should be promoted. It is also important to advise people on how to access these services.

Our findings suggest that providing opportunities for social activities and interaction for the hearing-impaired older adult is a viable intervention to promote social participation

**Table 2.** Hazard Ratios (HR) to Develop Dementia by Social Activities.

Hearingstatus	Explanatory variables	Categories	Crude Model		Model 1		Model 2	
			HR (95% CI)	p-value	HR (95% CI)	p-value	HR (95% CI)	p-value
Hearing impairment								
Sport group		>1 month	Reference		Reference		Reference	
		≤1 month	2.73 (2.02-3.69)	<0.01	2.17 (1.61-2.94)	<0.01	2.17 (1.53-3.08)	<0.01
Hobby group		>1 month	Reference		Reference		Reference	
		≤1 month	1.99 (1.63-2.44)	<0.01	1.70 (1.39-2.08)	<0.01	1.70 (1.34-2.17)	<0.01
Going out		>2 times a week	Reference		Reference		Reference	
		>1 times a month	1.78 (1.46-2.19)	0.02	1.28 (1.05-1.57)	0.02	1.29 (1.01-1.64)	0.04
		≤1 time a month	3.71 (2.75-5.00)	<0.01	2.32 (1.71-3.14)	<0.01	2.19 (1.51-3.17)	<0.01
Meeting with friends		>1 month	Reference		Reference		Reference	
		≤1 month	1.40 (1.20-1.63)	<0.01	1.35 (1.15-1.58)	<0.01	1.27 (1.06-1.53)	0.01
Leisure activities		Yes	Reference		Reference		Reference	
		No	1.78 (1.54-2.06)	<0.01	1.53 (1.32-1.78)	<0.01	1.57 (1.31-1.88)	<0.01
Normal hearing								
Sport group		>1 month	Reference		Reference		Reference	
		≤1 month	1.83 (1.67-2.01)	<0.01	1.52 (1.39-1.67)	<0.01	1.28 (1.16-1.42)	<0.01
Hobby group		>1 month	Reference		Reference		Reference	
		≤1 month	1.76 (1.64-1.89)	<0.01	1.55 (1.44-1.66)	<0.01	1.34 (1.23-1.45)	<0.01
Going out		>2 times a week	Reference		Reference		Reference	
		>1 times a month	2.44 (2.23-2.66)	<0.01	1.53 (1.40-1.67)	<0.01	1.44 (1.29-1.59)	<0.01
		≤1 time a month	5.07 (4.37-5.89)	<0.01	2.56 (2.20-2.98)	<0.01	2.09 (1.75-2.51)	<0.01
Meeting with friends		>1 month	Reference		Reference		Reference	
		≤1 month	1.48 (1.39-1.58)	<0.01	1.36 (1.28-1.45)	<0.01	1.24 (1.15-1.34)	<0.01
Leisure activities		Yes	Reference		Reference		Reference	
		No	2.14 (2.02-2.28)	<0.01	1.78 (1.68-1.89)	<0.01	1.55 (1.45-1.67)	<0.01

Model 1: adjusting for age, sex.

Model 2 further adjusting for self-rated health, activity of daily living, current smoking, and depressive symptoms as physical factors; living arrangement, equivalent income, education, and marital status as social factors.

HR: hazard ratio, CI: confidence interval.

among the elderly in Japan and may be effective in preventing the onset of dementia. Future research should investigate the types and details of social activities and interactions, such as sports and hobbies. This could contribute to the prevention of dementia in the hearing impaired.

### Strengths

The strengths of this study include its cohort design, which provides more convincing evidence than a cross-sectional design. Another strength is the large sample size of more

than 50 thousand total participants analyzed, and more than four thousand participants with HI. According to the large scale study, we revealed the relationship between social activities and the development of dementia in hearing-impaired older people.

### Limitations

This study had several limitations. First, self-administered questionnaires were used as the data collection tool, which may have led to information bias. The HI was assessed based on a questionnaire survey that uses responses to whether HI is being



treated. We were unable to exactly ascertain the severity of the HI in the participants, but we saw the HI level of participants as that which required treatment. There is a difference between the subjective assessment of hearing test results (Kamil et al., 2015) and the underestimation of HI (Uchida et al., 2003) in older adults. Second, information on the use of hearing aids and other devices that assist hearing function is missing. Among those receiving treatment for HI, it was not possible to discern who used hearing aids and other devices and who did not. Further research is needed to confirm these findings. Finally, this study did not consider factors in the community or the environment. Community-based intervention studies have shown that social capital effectively prevents the onset of disabilities and cognitive impairment (Hikichi et al., 2016; Hikichi et al., 2015). Community-level social capital has been associated with individual-level health outcomes (Tsuji et al., 2018; Tsuji et al., 2019; Fujihara et al., 2019). Urban areas, railroad tracks, airports, and highways are noisy, and the possibility of noise-induced HI cannot be ruled out (Chen et al., 2017). Environmental factors may also affect social activities, such as differences between urban and rural areas. In the future, it will be necessary to examine the differences between regions to determine how to address these issues at the community level.

In conclusion, this cohort study suggests that social interaction and social activities are related to cognitive function in people with HI. Maintaining opportunities for social communication and activities may help prevent dementia.

### Acknowledgments

This study used data from the JAGES. We are extremely grateful to all the study participants for their personal data. We would like to express our deepest gratitude to everyone who participated in the survey.

### Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study used data from the Japan Gerontological Evaluation Study (JAGES), which was supported by JSPS (Japan Society for the Promotion of Science) KAKENHI Grant Number (15H01972, 18H03047, 19H03860, 20H00557, 21K02001, 21K19635), Health Labour Sciences Research Grant (H28-Choju-Ippan-002), Japan Agency for Medical Research and Development (AMED) (17dk0110017, 18dk0110027, 18ls0110002, 18le0110009, 21dk0310108h0002, 22dk0310108h0003), the Research Funding for Longevity Sciences from National Center for Geriatrics and Gerontology (29-42, 21-20). Open Innovation Platform with Enterprises, Research Institute and Academia (OPERA, JPMJOP1831) from the Japan Science and Technology (JST).

### Authorship

Kaori Kojima (1) Substantial contributions to the conception or design of the work, or acquisition, analysis or interpretation of data for the work; (2) Drafting the work or revising it critically for important intellectual content; (3) Final approval of the version to be published; (4) Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. Meeting these criteria should provide each author with sufficient knowledge of and participation in the work that he or she can accept public responsibility for the report. Eisaku Okada (1) Substantial contributions to the conception or design of the work, or acquisition, analysis or interpretation of data for the work; (3) Final approval of the version to be published; (4) Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. Meeting these criteria should provide each author with sufficient knowledge of and participation in the work that he or she can accept public responsibility for the report. Toshiyuki Ojima (1) Substantial contributions to the conception or design of the work, or acquisition, analysis or interpretation of data for the work; (3) Final approval of the version to be published; (4) Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. Meeting these criteria should provide each author with sufficient knowledge of and participation in the work that he or she can accept public responsibility for the report. Jun Aida (2) Drafting the work or revising it critically for important intellectual content; (3) Final approval of the version to be published. Yoshimune Hiratsuka (2) Drafting the work or revising it critically for important intellectual content; (3) Final approval of the version to be published. Katsunori Kondo (2) Drafting the work or revising it critically for important intellectual content; (3) Final approval of the version to be published.

### ORCID iD

Kaori Kojima  <https://orcid.org/0000-0002-9593-5479>

### References

- Aida, J., Kondo, K., Hirai, H., Subramanian, S. V., Murata, C., Kondo, N., Ichida, Y., Shirai, K., & Osaka, K. (2011). Assessing the association between all-cause mortality and multiple aspects of individual social capital among the older Japanese. *BMC Public Health, 11*(1), 499. <https://doi.org/10.1186/1471-2458-11-499>
- Amieva, H., Ouvrard, C., Meillon, C., Rullier, L., & Dartigues, J. F. (2018). Death, depression, disability, and dementia associated with self-reported hearing problems: A 25-year study. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences, 73*(10), 1383–1389. <https://doi.org/10.1093/gerona/glx250>
- Barnes, L. L., Mendes de Leon, C. F., Wilson, R. S., Bienias, J. L., & Evans, D. A. (2004). Social resources and cognitive decline in a

- population of older African Americans and whites. *Neurology*, 63(12), 2322–2326. <https://doi.org/10.1212/01.wnl.0000147473.04043.b3>
- Bat-Chava, Y., Martin, D., & Kosciw, J. G. (2005). Longitudinal improvements in communication and socialization of deaf children with cochlear implants and hearing aids: Evidence from parental reports. *Journal of Child Psychology & Psychiatry*, 46(12), 1287–1296. <https://doi.org/10.1111/j.1469-7610.2005.01426.x>
- Bennett, D. A., Schneider, J. A., Arvanitakis, Z., Kelly, J. F., Aggarwal, N. T., Shah, R. C., & Wilson, R. S. (2006). Neuropathology of older persons without cognitive impairment from two community-based studies. *Neurology*, 66(12), 1837–1844. <https://doi.org/10.1212/01.wnl.0000219668.47116.e6>
- Berkman, L. F. (1995). The role of social relations in health promotion. *Psychosomatic Medicine*, 57(3), 245–254. <https://doi.org/10.1097/00006842-199505000-00006>
- Brice, P. J., & Strauss, G. (2016). Deaf adolescents in a hearing world: A review of factors affecting psychosocial adaptation. *Adolescent Health, Medicine and Therapeutics*, 7, 67–76. <https://doi.org/10.2147/AHMT.S60261>
- Brink, T. L., Yesavage, J. A., Lum, O., Heersema, P. H., Adey, M., & Rose, T. L. (1982). Screening tests for geriatric depression. *Clinical Gerontologist*, 1(1), 37–43. [https://doi.org/10.1300/J018v01n01\\_06](https://doi.org/10.1300/J018v01n01_06)
- Chen, H., Kwong, J. C., Copes, R., Tu, K., Villeneuve, P. J., van Donkelaar, A., Hystad, P., Martin, R. V., Murray, B. J., Jessiman, B., Wilton, A. S., Kopp, A., & Burnett, R. T. (2017). Living near major roads and the incidence of dementia, Parkinson's disease, and multiple sclerosis: A population-based cohort study. *Lancet (London, England)*, 389(10070), 718–726. [https://doi.org/10.1016/S0140-6736\(16\)32399-6](https://doi.org/10.1016/S0140-6736(16)32399-6)
- Daniele, A., Liguori, A., Bellomo, A., Greco, A., Panza, F., Lozupone, M., Sardone, R., Battista, P., Piccininni, M., Dibello, V., La Montagna, M., Stallone, R., Venezia, P., Liguori, A., Giannelli, G., Bellomo, A., Greco, A., Seripa, D., Quaranta, N., & Logroscino, G. (2018). Sensorial frailty: Age-related hearing loss and the risk of cognitive impairment and dementia in later life. *Therapeutic Advances in Chronic Disease*, 10, 2040622318811000. <https://doi.org/10.1177/2040622318811000>
- Dawes, P., Emsley, R., Cruickshanks, K. J., Moore, D. R., Fortnum, H., Edmondson-Jones, M., McCormack, A., & Munro, K. J. (2015). Hearing loss and cognition: The role of hearing AIDs, social isolation and depression. *PLoS One*, 10(3), Article e0119616. <https://doi.org/10.1371/journal.pone.0119616>
- Dori, D., Casale, G., Solerte, S. B., Fioravanti, M., Migliorati, G., Cuzzoni, G., & Ferrari, E. (1994). Chrono-neuroendocrinological aspects of physiological aging and senile dementia. *Chronobiologia*, 21(1–2), 121–126.
- Emmett, S. D., & Francis, H. W. (2015). The socioeconomic impact of hearing loss in US adults. *Otology & Neurotology*, 36(3), 545–550. <https://doi.org/10.1097/mao.0000000000000562>
- Fratiglioni, L., Wang, H. X., Ericsson, K., Maytan, M., & Winblad, B. (2000). Influence of social network on occurrence of dementia: A community-based longitudinal study. *Lancet (London, England)*, 355(9212), 1315–1319. [https://doi.org/10.1016/S0140-6736\(00\)02113-9](https://doi.org/10.1016/S0140-6736(00)02113-9)
- Fujihara, S., Tsuji, T., Miyaguni, Y., Aida, J., Saito, M., Koyama, S., & Kondo, K. (2019). Does community-level social capital predict decline in instrumental activities of daily living? A JAGES prospective cohort study. *International Journal of Environmental Research and Public Health*, 16(5), 828. <https://doi.org/10.3390/ijerph16050828>
- Hikichi, H., Kondo, N., Kondo, K., Aida, J., Takeda, T., & Kawachi, I. (2015). Effect of a community intervention programme promoting social interactions on functional disability prevention for older adults: Propensity score matching and instrumental variable analyses, JAGES Taketoyo study. *Journal of Epidemiology and Community Health*, 69(9), 905–910. <https://doi.org/10.1136/jech-2014-205345>
- Hikichi, H., Kondo, K., Takeda, T., & Kawachi, I. (2016). Social interaction and cognitive decline: Results of a 7-year community intervention. *Alzheimer's & Dementia (New York, N. Y.)*, 3(1), 23–32. <https://doi.org/10.1016/j.trci.2016.11.003>
- Hisano, S. (2009). The relationship between revised hasegawa dementia scale (HDS-R), mini-mental state examination (MMSE) and bed-fast scale, dementia scale. *Japanese Journal of Geriatric Psychiatry*, 20(8), 883–891. [in Japanese].
- Holtzman, R. E., Rebok, G. W., Saczynski, J. S., Kouzis, A. C., Wilcox Doyle, K., & Eaton, W. W. (2004). Social network characteristics and cognition in middle-aged and older adults. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 59(6), P278–P284. <https://doi.org/10.1093/geronb/59.6.p278>
- Ives, D. G., Bonino, P., Traven, N. D., & Kuller, L. H. (1995). Characteristics and comorbidities of rural older adults with hearing impairment. *Journal of the American Geriatrics Society*, 43(7), 803–806. <https://doi.org/10.1111/j.1532-5415.1995.tb07056.x>
- Kamil, R. J., Genther, D. J., & Lin, F. R. (2015). Factors associated with the accuracy of subjective assessments of hearing impairment. *Ear and Hearing*, 36(1), 164–167. <https://doi.org/10.1097/AUD.0000000000000075>
- Kanamori, S., Kai, Y., Aida, J., Kondo, K., Kawachi, I., Hirai, H., Shirai, K., Ishikawa, Y., & Suzuki, K., JAGES Group. (2014). Social participation and the prevention of functional disability in older Japanese: The JAGES cohort study. *PLoS One*, 9(6), Article e99638. <https://doi.org/10.1371/journal.pone.0099638>
- Kawachi, I., Colditz, G. A., Ascherio, A. E. B., Rimm, E. B., Giovannucci, E., Stampfer, M. J., & Willett, W.C. (1996). A prospective study of social networks in relation to total mortality and cardiovascular disease in men in the USA. *Journal of Epidemiology and Community Health*, 50(3), 245–251. <https://doi.org/10.1136/jech.50.3.245>
- Knutson, J. F., & Lansing, C. R. (1990). The relationship between communication problems and psychological difficulties in persons with profound acquired hearing loss. *Journal of Speech & Hearing Disorders*, 55(4), 656–664. <https://doi.org/10.1044/jshd.5504.656>

- Kochkin, S. (2007). *The impact of untreated hearing loss on household income*. Better Hearing Institute.
- Levasseur, M., Richard, L., Gauvin, L., & Raymond, E. (2010). Inventory and analysis of definitions of social participation found in the aging literature: Proposed taxonomy of social activities. *Social Science & Medicine* (1982), 71(12), 2141–2149. <https://doi.org/10.1016/j.socscimed.2010.09.041>
- Livingston, G., Sommerlad, A., Orgeta, V., Costafreda, S. G., Huntley, J., Ames, D., Ballard, C., Banerjee, S., Burns, A., Cohen-Mansfield, J., Cooper, C., Fox, N., Gitlin, L. N., Howard, R., Kales, H. C., Larson, E. B., Ritchie, K., Rockwood, K., Sampson, E. L., & Mukadam, N. (2017). Dementia prevention, intervention, and care. *Lancet (London, England)*, 390(10113), 2673–2734. [https://doi.org/10.1016/S0140-6736\(17\)31363-6](https://doi.org/10.1016/S0140-6736(17)31363-6)
- Li, C. M., Zhang, X., Hoffman, H. J., Cotch, M. F., Themann, C. L., & Wilson, M. R. (2014). Hearing impairment associated with depression in US adults, national health and nutrition examination survey 2005-2010. *JAMA Otolaryngology Head & Neck Surgery*, 140(4), 293–302. <https://doi.org/10.1001/jamaoto.2014.42>
- Michalowsky, B., Hoffmann, W., & Kostev, K. (2019). Association between hearing and vision impairment and risk of dementia: Results of a case—control study based on secondary data. *Frontiers in Aging Neuroscience*, 11, 363. <https://doi.org/10.3389/fnagi.2019.00363>
- Moog, J. S., Geers, A. E., Gustus, C. H., & Brenner, C. A. (2011). Psychosocial adjustment in adolescents who have used cochlear implants since preschool. *Ear and Hearing*, 32(Suppl 1), 75S–83S. <https://doi.org/10.1097/AUD.0b013e3182014c76>
- Murata, C., Saito, T., Tsuji, T., Saito, M., & Kondo, K. (2017). A 10-year follow-up study of social ties and functional health among the Old: The AGES Project. *International Journal of Environmental Research and Public Health*, 14(7), 717. <https://doi.org/10.3390/ijerph14070717>
- Nemoto, Y., Saito, T., Kanamori, S., Arao, T., Tsuji, T., Kikuchi, H., Maruo, K., Arao, T., Kondo, K., Yuta, N., Tami, S., Satoru, K., Taishi, T., Kokoro, S., Hiroyuki, K., Shirai, K., Maruo, K., Arao, R., & Kondo, K. (2017). An additive effect of leading role in the organization between social participation and dementia onset among Japanese older adults: The AGES cohort study. *BMC Geriatrics* 17(1), 297. <https://doi.org/10.1186/s12877-017-0688-9>
- Noda, H., Yamagishi, K., Ikeda, A., Asada, T., & Iso, H. (2018). Identification of dementia using standard clinical assessments by primary care physicians in Japan. *Geriatrics & Gerontology International*, 18(5), 738–744. <https://doi.org/10.1111/ggi.13243>
- Ogawa, T., Uchida, Y., Nishita, Y., Tange, C., Sugiura, S., Ueda, H., Nakada, T., Suzuki, H., Otsuka, R., Ando, F., & Shimokata, H. (2019). Hearing-impaired elderly people have smaller social networks: A population-based aging study. *Archives of Gerontology and Geriatrics*, 83, 75–80. <https://doi.org/10.1016/j.archger.2019.03.004>
- Panza, F., Solfrizzi, V., & Logroscino, G. (2015). Age-related hearing impairment—a risk factor and frailty marker for dementia and AD. *Nature Reviews Neurology*, 11(3), 166–175. <https://doi.org/10.1038/nrneuro.2015.12>
- Pérès, K., Helmer, C., Amieva, H., Orgogozo, J. M., Rouch, I., Dartigues, J. F., & Barberger-Gateau, P. (2008). Natural history of decline in instrumental activities of daily living performance over the 10 years preceding the clinical diagnosis of dementia: A prospective population-based study. *Journal of the American Geriatrics Society*, 56(1), 37–44. <https://doi.org/10.1111/j.1532-5415.2007.01499.x>
- Putnam, R. D. (1995). Bowling alone: America's declining social capital. *Journal of Democracy*, 6(1), 65–78. <https://doi.org/10.1353/jod.1995.0002>
- Saczynski, J. S., Pfeifer, L. A., Masaki, K., Korf, E. S., Laurin, D., White, L., & Launer, L. J. (2006). The effect of social engagement on incident dementia: The Honolulu-Asia aging study. *American Journal of Epidemiology*, 163(5), 433–440. <https://doi.org/10.1093/aje/kwj061>
- Saito, M., Aida, J., Kondo, N., Saito, J., Kato, H., Ota, Y., Amemiya, A., & Kondo, K. (2019). Reduced long-term care cost by social participation among older Japanese adults: A prospective follow-up study in JAGES. *BMJ Open*, 9(3), Article e024439. <https://doi.org/10.1136/bmjopen-2018-024439>
- Servidoni, A. B., & Conterno, L. O. (2018). Hearing loss in the elderly: Is the hearing handicap inventory for the elderly - screening version effective in diagnosis when compared to the audiometric test? *International Archives of Otorhinolaryngology*, 22(1), 1–8. <https://doi.org/10.1055/s-0037-1601427>
- Strawbridge, W. J., Wallhagen, M. I., Shema, S. J., & Kaplan, G. A. (2000). Negative consequences of hearing impairment in old age: A longitudinal analysis. *The Gerontologist*, 40(3), 320–326. <https://doi.org/10.1093/geront/40.3.320>
- Sugawara, N., Sasaki, A., Yasui-Furukori, N., Kakehata, S., Umeda, T., Namba, A., Nakaji, S., Shinkawa, H., & Kaneko, S. (2011). Hearing impairment and cognitive function among a community-dwelling population in Japan. *Annals of General Psychiatry*, 10(1), 27. <https://doi.org/10.1186/1744-859X-10-27>
- Tsuji, T., Kanamori, S., Miyaguni, Y., Hanazato, M., & Kondo, K. (2019). Community-level sports group participation and the risk of cognitive impairment. *Medicine and Science in Sports and Exercise*, 51(11), 2217–2223. <https://doi.org/10.1249/MSS.0000000000002050>
- Tsuji, T., Y., Miyaguni, Y., Kanamori, S., Hanazato, M., & Kondo, K. (2018). Community-level Sports Group Participation and Older Individuals' Depressive Symptoms. *Medicine and science in sports and exercise*, 50(6), 1199–1205. <https://doi.org/10.1249/MSS.0000000000001541>
- Uchida, Y., Nakashima, T., Ando, F., Niino, N., & Shimokata, H. (2003). Prevalence of self-perceived auditory problems and their relation to audiometric thresholds in a middle-aged to elderly population. *Acta Oto-Laryngologica*, 123(5), 618–626. <https://doi.org/10.1080/00016480310001448>
- Uchida, Y., Sugiura, S., Nishita, Y., Saji, N., Sone, M., & Ueda, H. (2019). Age-related hearing loss and cognitive decline - The potential mechanisms linking the two. *Auris, Nasus, Larynx*, 46(1), 1–9. <https://doi.org/10.1016/j.anl.2018.08.010>

- Uchino, B. N. (2006) Social support and health: A review of physiological processes potentially underlying links to disease outcomes. *Journal of Behavioral Medicine*, 29(4), 377–387. <https://doi.org/10.1007/s10865-006-9056-5>
- Wayne, R. V., & Johnsrude, I. S. (2015). A review of causal mechanisms underlying the link between age-related hearing loss and cognitive decline. *Ageing Research Reviews*, 23, 154166, <https://doi.org/10.1016/j.arr.2015.06.002>
- Willie-Tyndale, D., Holder-Nevins, D., Mitchell-Fearon, K., James, K., Laws, H., Waldron, N. K., & Eldemire-Shearer, D. (2016). Participation in social activities and the association with socio-demographic and health-related factors among community-dwelling older adults in Jamaica. *Journal of Cross-Cultural Gerontology*, 31(4), 427–447. <https://doi.org/10.1007/s10823-016-9297-x>
- Yamada, M., Nishiwaki, Y., Michikawa, T., & Takebayashi, T. (2011). Impact of hearing difficulty on dependence in activities of daily living (ADL) and mortality: A 3-year cohort study of community-dwelling Japanese older adults. *Archives of Gerontology and Geriatrics*, 52(3), 245–249. <https://doi.org/10.1016/j.archger.2010.04.023>