

Efficacy of stress measurements using salivary ultra-weak chemiluminescence

メタデータ	言語: eng 出版者: 日本職業・災害医学会 公開日: 2019-07-22 キーワード (Ja): キーワード (En): 作成者: 中村, 正光 メールアドレス: 所属:
URL	http://hdl.handle.net/10271/00003573

Efficacy of stress measurements using salivary ultra-weak chemiluminescence

Masamitsu Nakamura^{1,2)}, Kazuo Umemura³⁾, Mitsuo Hiramatsu⁴⁾, Hiroko Oishi⁵⁾, Masato Maekawa⁶⁾

¹⁾ Graduate School of Medicine, Hamamatsu University School of Medicine

²⁾ Department of Laboratory, Hamamatsu Rosai Hospital

³⁾ Department of Pharmacology, Hamamatsu University School of Medicine

⁴⁾ Central Research Laboratory, Hamamatsu Photonics K.K.,

⁵⁾ Department of Endocrinology, Hamamatsu Rosai Hospital

⁶⁾ Department of Laboratory Medicine, Hamamatsu University School of Medicine

Abstract

The incidence of stress-induced psychiatric disorders has increased rapidly in recent years. Screening for stress markers enables early detection and treatment of conditions in which stress responses affect health. Salivary ultra-weak chemiluminescence (UCL) can detect peroxidase and thiocyanate ions, which are secreted and increased in saliva under stress. Forty-four patients with diabetes mellitus who were admitted for diabetic education because of their high prevalence of depression and 14 healthy hospital staff members were included. Level variations of stress-related substances in saliva were measured by UCL. Simultaneously, salivary levels of cortisol and chromograninA (CgA) were measured, and the short form of the Profile of Mood States questionnaire was administered. Chemiluminescence intensity correlated with increases in depression-dejection subscale scores, and had a positive and negative correlation with CgA and cortisol levels, respectively. The hospital staff and patients differed significantly in chemiluminescence intensities, cortisol levels, and depression-dejection scores. In conclusion, UCL intensity was correlated with depression scores and levels of classical stress markers, with significant differences between hospital staff and patients. UCL intensity may thus allow the measurement of increases in psychological stress. Furthermore, because saliva absorbed by filter paper can be used, samples can easily be collected and analyzed.

—Keywords—

ultra-weak chemiluminescence, Profile of Mood States, stress marker

Introduction

In recent years, the incidence of stress-induced psychiatric disorders, such as depression, has increased

rapidly. At present, stress is mainly measured and assessed using psychological questionnaires, which need to be administered and evaluated by physicians trained in psychology or by psychologists. Meanwhile, although measurements of biomarkers such as cortisol, amylase, chromogranin A (CgA), and immunoglobulin A (IgA),¹⁾⁻⁶⁾ in blood, urine, and saliva have been reported to be useful, these biomarkers have not been studied for long enough to be adequate and quantifiable markers. Stress markers that are easy to measure and highly specific seem to be needed. Salivary ultra-weak chemiluminescence (UCL) can detect peroxidase and thiocyanate ions, which are secreted and increased in saliva under stress.⁷⁾⁻⁹⁾ It has been reported that salivary UCL intensity significantly increased in patients with severe diseases, such as lung cancer,⁸⁾ and in people under stress caused by mental workload in the Kraepelin test.⁹⁾ Salivary UCL intensity can be measured in saliva absorbed by filter paper, which gives this marker an advantage in that saliva samples can be collected easily and repeatedly.⁷⁾⁻⁹⁾

This study targeted patients admitted to a hospital for diabetic education. According to a statistical survey conducted by the Japanese Ministry of Health, Labour and Welfare in 2011,¹⁰⁾ diabetes mellitus affects approximately 10.67 million of the Japanese adult population of approximately 95.34 million, and its prevalence reached 11.2% in 2011. While diabetes mellitus causes cardiovascular complications, the prevalence of depression is also reported to be as high as 14% in patients with diabetes mellitus.¹⁰⁾ Those with concomitant depression are unable to exercise glycemic self-control, which leads to a vicious cycle. When we previously administered the Profile of Mood States (POMS) questionnaire to patients admitted for diabetic education, psychological stress was observed in 11 of 16 surveyed patients (unpublished data). This observation indicates that patients with severe psychological stress are likely to be encountered in such a patient group. Thus, by targeting patients admitted for diabetic education, this study aimed to compare salivary UCL to a conventional psychological questionnaire as well as to measurements of salivary CgA and cortisol, to determine whether salivary UCL intensity could be a useful stress marker in the future.

Experimental

1. Subjects and methods

In this study, salivary UCL, the POMS questionnaire, and conventional measurements of salivary CgA and cortisol were analyzed and compared to determine the usefulness of salivary UCL for stress measurements. This study included 44 patients admitted for diabetic education (31 men and 13 women) and 14 hospital staff members without diabetes mellitus (seven males and seven females). Prior to this study, the experimental protocol was approved by the ethics committee of Hamamatsu Rosai Hospital, and informed consent for participation in the study was obtained from all study participants.

2. Sample collection

During the diabetic education sessions, the POMS questionnaire was administered, and saliva samples were collected. In these sessions which were held between 15:00 and 16:00 p.m. before the evening meal, the patients were examined after receiving instructions for diabetes treatment while seated for approximately 30 minutes, to avoid any effects of exercise. The hospital staff members were also examined before meal and under resting conditions. Study participants who were willing to repeatedly participate in this

study underwent several examinations on different days.

Salivary ultra-weak chemiluminescence (UCL) and levels of salivary chromogranin A (CgA) and salivary cortisol were measured in the samples from the study participants. Filter paper was used for salivary UCL, and the Saliva Collection Aid (SCA; Salimetrics LLC, USA) was used for measurements of CgA and cortisol. The filter paper (Nobuto Blood-Sampling Paper Strip type I) was placed in the mouth to absorb saliva for 60 seconds. Each filter strip that had absorbed saliva was placed in a zip-seal plastic bag, frozen, and stored at -80°C until measurement.

3. Measurements

Salivary UCL intensity was measured with a Photon Counter C767 (Hamamatsu Photonics K.K., Hamamatsu, Japan).⁷⁾ The cryopreserved filter paper that had absorbed saliva was brought to room temperature under running water. After 1 mL of a gallate solution adjusted to a concentration of 3 mM (gallate 0.282 g + distilled water 500 mL) and the filter paper were placed in the measurement cell, the measurement started. When a 3% hydrogen peroxide solution was added 20 seconds later, chemiluminescence started. The mean intensity of chemiluminescence measured for 100 seconds was used for analysis. (Fig 1)

Fig 1




Fig.1. Salivary UCL levels. After 3% (w/v) H₂O₂ solution was added to the saliva preparation, photo counts were conducted. The total number of counts for 100 seconds was defined as the level of UCL.

Salivary CgA levels were measured by enzyme immunoassay using a microplate reader (SPECTRAmax(R) PLUS (384); Molecular Devices Inc., USA) and calibrated to protein equivalents (expressed as pmol/mg). Salivary cortisol levels were measured by solid-phase radioimmunoassay using a γ counter (Aloka ARC-950; Hitachi-Aloka Medical, Tokyo, Japan) and expressed as $\mu\text{g/dL}$.

Results

Table 1 shows the number of saliva samples collected from the study participants according to the study parameters (UCL, CgA, and cortisol), the number of times when the POMS questionnaire was administered, and the mean age of the study participants.

For measurements of salivary UCL intensity, 83 saliva samples (55 from male and 28 from female participants) were collected overall, and the mean age at the time of this study was 55 ± 15 years (55 ± 14 vs. 55 ± 16 for male/female participants). From the patients admitted for diabetic education, 49 saliva samples (37 vs. 12) were collected, and the mean age at the time of this study was 57 ± 14 years (57 ± 14 vs. 59 ± 13). From the hospital staff members, 34 saliva samples (18 from male and 16 from female participants) were collected, and the mean age at the time of this study was 42 ± 13 years (47 ± 11 vs. 36 ± 13). The survey and measurement results of individual participants are shown in detail in the supplementary tables.

The salivary UCL intensity measured in 83 saliva samples widely varied from 149 to 20000 average counts/100 seconds.

Table 1

Sample size according to the study parameters and average age

	All participants			Hospital staff			Patients		
	total	male	female	total	male	female	total	male	female
UCL	83	55	28	34	18	16	49	37	12
CgA	70	48	22	33	17	16	37	31	6
Cortisol	70	50	20	33	18	15	37	32	5
POMS	82	55	27	33	18	15	49	37	12
Average Age	55 ± 15	55 ± 14	55 ± 16	42 ± 13	47 ± 11	36 ± 13	57 ± 14	57 ± 14	59 ± 13

1. POMS questionnaire

For the POMS questionnaire, responses were converted to numerical scores. In this study, the vigor POMS subscale scores (healthy range: ≥ 8 for men and ≥ 9 for women) decreased in 32 of the 58 study participants (25 patients admitted for diabetic education and seven hospital staff members). Meanwhile, the POMS scores increased on the tension-anxiety subscale (healthy range: ≥ 11 for men and ≥ 12 for women) in 11 participants (eight patients and three hospital staff members), on the anger-hostility subscale (healthy range: ≤ 9 for both men and women) in nine participants (nine patients), on the fatigue subscale (healthy range: ≤ 12 for men and ≤ 13 for women) in eight participants (seven patients and one hospital staff member), and on the confusion subscale (healthy range: ≤ 9 for both men and women) in 12 participants (nine patients and three hospital staff members). Depression (dejection) (healthy range of the POMS subscale score: ≤ 8 for men and ≤ 9 for women) was observed in 10 participants (eight patients and two hospital staff members). In some surveyed patients whose profiles indicated that they were treated for depression, suffered from insomnia, or were under excessive stress from working at night for a long period, the POMS scores increased on the depression-

dejection and tension-anxiety subscales and decreased on the vigor subscale. When we examined whether salivary UCL intensity measured in 82 saliva samples was correlated with increases and decreases in the depression-dejection POMS subscale scores, a correlation was observed between intensity and scores ($p < 0.05$, $r = 0.24$). (Fig.2)




Fig.2. POMS questionnaire. There was a significant correlation between UCL intensity and the depression-dejection POMS subscale scores ($p < 0.05$, $r = 0.24$; 82 saliva samples).

2. Comparison between hospital staff and patients

The comparison between the hospital staff and the patient groups revealed significant differences in salivary UCL intensity ($p < 0.05$), cortisol levels ($p < 0.05$), and depression-dejection POMS subscale scores ($0.05 < p < 0.1$). (Table 2)

Table 2

Hospital staff crowd and Patients

	Hospital staff	Patients	P-value
UCL (avarege counts/100 s)	490.5 ± 270.9	3330.9 ± 5445.2	p < 0.05
CgA (pml/ml)	4.2 ± 3.6	5.5 ± 3.2	ns
Cortisol (IU/L)	0.3 ± 0.3	0.2 ± 0.1	p < 0.05
Depression-Dejection (score)	3.1 ± 3.6	4.5 ± 3.8	0.05 < p < 0.1

3. Chromogranin A

CgA levels varied from 0.5 to 14.6 pmol/mg of protein. Although no correlation was observed with the depression-dejection POMS subscale scores, CgA levels were correlated with salivary UCL intensity measured in 70 saliva samples ($p < 0.05$, $r = 0.23$). A Spearman rank correlation analysis also showed a positive correlation ($r_s = 0.388$). (Fig. 3)

Fig3




Fig.3. Chromogranin A. CgA levels were correlated with salivary UCL intensity measured in 70 saliva samples ($p < 0.05$, $r = 0.23$).

4. Cortisol

Cortisol levels ranged from 0.06 to 1.20 $\mu\text{g/dL}$. When compared between the hospital staff and the patient groups, the levels were correlated with the depression-dejection POMS subscale scores ($p < 0.05$). The levels also showed a significant difference between the groups ($p < 0.05$). When we examined whether the cortisol levels were correlated with salivary UCL intensity measured in 70 saliva samples, a negative correlation was observed ($p < 0.05$). A Spearman rank correlation analysis also showed a negative correlation ($r_s = -0.278$). (Fig.4)

Fig4




Fig.4. Cortisol levels. There was a negative correlation between UCL intensity and cortisol levels in 70 saliva samples ($p < 0.05$).

Discussion

In this study, we examined whether salivary UCL intensity is an effective stress marker. The study included patients admitted to a hospital for diabetic education. Simultaneously, the POMS questionnaire, which is most commonly used for assessing psychological stress, was administered to the patients. In some surveyed

patients whose profiles indicated that they were treated for depression, suffered insomnia, or were under excessive stress from working at night for a long period, the POMS scores increased on the depression-dejection and tension-anxiety subscales, and decreased on the vigor subscale. The salivary UCL intensity measured in 82 saliva samples was correlated with increases in the POMS scores, particularly on the depression-dejection subscale. These findings suggest UCL intensity as a candidate for a reliable stress marker.

Salivary level of CgA, which is a classical stress marker, was simultaneously measured. A Spearman rank correlation analysis of the measurement and salivary UCL intensity revealed a positive correlation between salivary CgA levels and UCL intensity. Salivary CgA levels increase under stress and mental load, such as tension and fear, in the early stages. These findings thus suggest that UCL intensity can detect stress as well as the classical stress marker CgA.

Cortisol levels tend to increase under intensive and persistent physical stress or chronic stress,¹¹⁾ but have been reported to be normal or low under acute stress.^{12),13)} This study revealed a negative correlation between salivary cortisol levels and UCL intensity. When we had previously measured UCL intensity before and after mental workload in the Kraepelin test,⁹⁾ the acute stress caused increases in UCL intensity levels. UCL levels are considered to increase during acute stress. Previous results and this study thus suggest that salivary UCL intensity increases with acute stress.

In this study, measurements obtained from patients with marked gingivitis varied, maybe due to bleeding in the oral cavity. Therefore, before saliva samples are collected for salivary UCL, gargling is necessary to reduce the influence of bleeding and other factors in the oral cavity. However, measurements of salivary UCL intensity widely varied from 149 to 200000 average counts/100 seconds in this study, and we were unable to explain why they increased to such a high level.

The collection of saliva samples from patients with diabetes mellitus is often difficult in patients whose salivary secretion is low. Saliva samples for UCL can however be easily collected, even in patients with low salivary secretion, because salivary UCL intensity can be measured in saliva absorbed by filter paper.

In conclusion, we found that salivary UCL intensity is correlated with depression-dejection POMS subscale scores; it is also positively correlated with salivary CgA levels and negatively correlated with salivary cortisol levels. These findings suggest that UCL intensity can be used as an acute stress marker. Furthermore, because salivary UCL intensity can be measured in saliva absorbed by filter paper, saliva samples can be collected easily.

Abbreviations

ultra-weak chemiluminescence (UCL); chromogranin A (CgA); Profile of Mood States (POMS); Saliva Collection Aid (SCA)

Acknowledgement

None

References

- 1) Vivian Ng, B.Sc. (Hons), David Koh, et al. :Salivary Biomarkers Associated with Academic Assessment Stress Among Dental Undergraduates. *Journal of Dental Education*. 67 2003, pp1091-1094
- 2) Koichi Takatsuji, Yoshie Sugimoto, Shoko Ishizaki, et al. :The effects of examination stress on salivary cortisol, immunoglobulin A, and chromogranin A in nursing students. *Biomedical Research* 29 2008, pp221-224
- 3) Jana Strahler, Nadine Skoluda, Mattes B kappert, et al. :Simultaneous measurement of salivary cortisol and alpha-amylase: Application and recommendations. *Neuroscience and Biobehavioral Reviews* 83 2017 pp657-677
- 4) Hitoshi Abekura, Masahiro Tsuboi, Tomohisa Okura, et al :Association between sleep bruxism and stress sensitivity in an experimental psychological stress task. *Biomedical Research* 32 2011 pp395-399
- 5) Tamaki Matsumoto, Hiroyuki Asakura, Tatsuya Hayashi. :Effects of Olfactory Stimulation from the Fragrance of the Japanese Citrus Fruit Yuzu (*Citrus junos* Sieb. Ex Tanaka) on Mood States and Salivary Chromogranin A as an Endocrinologic Stress Maker. *The journal of alternative and complementary medicine* 20 2013 pp500-506
- 6) Tamaki Matsumoto, Hiroyuki Asakura, Tatsuya Hayashi. :Increased salivary chromograninA in women with severe negative mood states in the premenstrual phase. *J Psychosom obstet gynaecol*. 33 2012 pp120-128
- 7) Kuniaki Takagi, Tsuyoshi Hasunuma, Nobuhiro Goi, et al, Short Communication: The effects of performance anxiety on salivary ultra-weak chemiluminescence. *Stress and Health* 21 2005 pp263-268
- 8) Mitsuo Hiramatsu, Kingo Chida, Dai Hashimoto, et al, Possibility of using salivary ultra-weak chemiluminescence as a biomarker for feelings of anxiety in hospital settings. *Luminescence* 25 2015 pp1395-1399
- 9) Nobuhiro Goi, Kuniaki Takagi, Yuuko Hirai, et al. Effect of Psychologic Stress on Peroxidase and Thiocyanate Levels in Human Saliva Detected by Ultraweak *Journal of Health Science*.53 2007 pp161-169
- 10) Nigel Unwin, David Whiting, Leonor Guariguata, et al. *Diabetes Atlas*. 5 th ed. International Diabetes Federation, Brussels, Belgium, 2011
- 11) Shusaku Nomura, Hideki Ohara, Tsutomu Kamei, Effect of the relief from chronic stress during graduation examination on salivary biomarkers. *Transactions of the Japan Society of Kasai Engineering* 8 2009 pp481-487
- 12) Helane Wahbeh, Barry S. Oken, Salivary Cortisol lower in posttraumatic stress disorder. *J Trauma Stress* 26 2013 pp1-15
- 13) Dawn M. Jonshon, Douglas L. Deahanty, et al. The cortisol awakening response as function of PTSD severity and abuse chronicity shelter battered women. *J Anxiety Disord* 2008 pp793-800

Reprint request:

Masamitsu Nakamura

Department of Laboratory, Hamamatsu Rosai Hospital, Shogencho 25, Higashi-ku, Hamamatsu 430-8525, Japan

別刷請求先 〒430-8525 静岡県浜松市東区将監町 25 浜松労災病院 中央検査部 中村正光