



Differences in fractional anisotropy between the patients with schizophrenia and healthy comparison subjects

メタデータ	<p>言語: English</p> <p>出版者:</p> <p>公開日: 2021-01-05</p> <p>キーワード (Ja):</p> <p>キーワード (En):</p> <p>作成者: Koshiyama, Daisuke, Fukunaga, Masaki, Okada, Naohiro, Morita, Kentaro, Nemoto, Kiyotaka, Usui, Kaori, Yamamori, Hidenaga, Yasuda, Yuka, Fujimoto, Michiko, Kudo, Noriko, Azechi, Hirotugu, Watanabe, Yoshiyuki, Hashimoto, Naoki, Narita, Hisashi, Kusumi, Ichiro, Ohi, Kazutaka, Shimada, Takamitsu, Kataoka, Yuzuru, Yamamoto, Maeri, Ozaki, Norio, Okada, Go, Okamoto, Yasumasa, Harada, Kenichiro, Matsuo, Koji, Yamasue, Hidenori, Abe, Osamu, Hashimoto, Ryuichiro, Takahashi, Tsutomu, Hori, Tomoki, Nakataki, Masahito, Onitsuka, Toshiaki, Holleran, Laurena, Jahanshad, Neda, van Erp, G. M. Theo, Turner, Jessica, Donohoe, Gary, Thompson, M. Paul, Kasai, Kiyoto, Hashimoto, Ryota, COCORO</p> <p>メールアドレス:</p> <p>所属:</p>
-------	--

Figure 1A Koshiyama et al.

