

# Liganded T3 receptor $\beta$ 2 inhibits the positive feedback autoregulation of the gene for GATA2, a transcription factor critical for thyrotropin production

メタデータ	言語: English 出版者: PLOS (Public Library of Science) 公開日: 2022-11-28 キーワード (Ja): キーワード (En): 作成者: 平原, 直子 メールアドレス: 所属:
URL	<a href="http://hdl.handle.net/10271/00004209">http://hdl.handle.net/10271/00004209</a>

This work is licensed under a Creative Commons Attribution 4.0 International License.



## RESEARCH ARTICLE


# Liganded T3 receptor $\beta 2$ inhibits the positive feedback autoregulation of the gene for GATA2, a transcription factor critical for thyrotropin production

Naoko Hirahara<sup>1</sup>, Hiroko Misawa Nakamura<sup>2</sup>, Shigekazu Sasaki<sup>1,2\*</sup>, Akio Matsushita<sup>2</sup>, Kenji Ohba<sup>3</sup>, Go Kuroda<sup>2</sup>, Yuki Sakai<sup>2</sup>, Shinsuke Shinkai<sup>2</sup>, Hiroshi Haeno<sup>4</sup>, Takuhiro Nishio<sup>5</sup>, Shuichi Yoshida<sup>5</sup>, Yutaka Oki<sup>6</sup>, Takafumi Suda<sup>2</sup>

**1** Division of Endocrinology and Metabolism, Department of Internal medicine, Japanese Red Cross Shizuoka Hospital, Shizuoka, Shizuoka, Japan, **2** Second Division, Department of Internal Medicine, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan, **3** Medical Education Center, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan, **4** Department of Computational Biology and Medical Sciences, Graduate School of Frontier Sciences, The University of Tokyo Kashiwa, Kashiwa, Chiba, Japan, **5** Department of Integrated Human Sciences, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan, **6** Department of Family and Community Medicine, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka, Japan

✉ These authors contributed equally to this work.

\* [sasakis@hama-med.ac.jp](mailto:sasakis@hama-med.ac.jp)



## OPEN ACCESS

**Citation:** Hirahara N, Nakamura HM, Sasaki S, Matsushita A, Ohba K, Kuroda G, et al. (2020) Liganded T3 receptor  $\beta 2$  inhibits the positive feedback autoregulation of the gene for GATA2, a transcription factor critical for thyrotropin production. *PLoS ONE* 15(1): e0227646. <https://doi.org/10.1371/journal.pone.0227646>

**Editor:** Hiroyoshi Ariga, Hokkaido Daigaku, JAPAN

**Received:** August 14, 2019

**Accepted:** December 23, 2019

**Published:** January 15, 2020

**Peer Review History:** PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0227646>

**Copyright:** © 2020 Hirahara et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its Supporting Information files.

## Abstract

The serum concentration of thyrotropin (thyroid stimulating hormone, TSH) is drastically reduced by small increase in the levels of thyroid hormones (T3 and its prohormone, T4); however, the mechanism underlying this relationship is unknown. TSH consists of the chorionic gonadotropin  $\alpha$  (CGA) and the  $\beta$  chain (TSH $\beta$ ). The expression of both peptides is induced by the transcription factor GATA2, a determinant of the thyrotroph and gonadotroph differentiation in the pituitary. We previously reported that the liganded T3 receptor (TR) inhibits transactivation activity of GATA2 via a tethering mechanism and proposed that this mechanism, but not binding of TR with a negative T3-responsive element, is the basis for the T3-dependent inhibition of the TSH $\beta$  and CGA genes. Multiple GATA-responsive elements (GATA-REs) also exist within the GATA2 gene itself and mediate the positive feedback autoregulation of this gene. To elucidate the effect of T3 on this non-linear regulation, we fused the GATA-REs at -3.9 kb or +9.5 kb of the GATA2 gene with the chloramphenicol acetyltransferase reporter gene harbored in its 1S-promoter. These constructs were co-transfected with the expression plasmids for GATA2 and the pituitary specific TR, TR $\beta 2$ , into kidney-derived CV1 cells. We found that liganded TR $\beta 2$  represses the GATA2-induced transactivation of these reporter genes. Multi-dimensional input function theory revealed that liganded TR $\beta 2$  functions as a classical transcriptional repressor. Then, we investigated the effect of T3 on the endogenous expression of GATA2 protein and mRNA in the gonadotroph-derived L $\beta$ T2 cells. In this cell line, T3 reduced GATA2 protein independently of the ubiquitin proteasome system. GATA2 mRNA was drastically suppressed by T3, the concentration of which corresponds to moderate hypothyroidism and euthyroidism. These results

**Funding:** This work was supported in part by a Grant-in Aid for Scientific Research to S. S. from the Ministry of Education, Culture, Sports, Science and Technology of Japan (grant number 24590689). The sponsors or funders play no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. <https://www.jsps.go.jp/j-grantsinaid/>

**Competing interests:** The authors have declared that no competing interests exist.

suggest that liganded TR $\beta$ 2 inhibits the positive feedback autoregulation of the GATA2 gene; moreover this mechanism plays an important role in the potent reduction of TSH production by T3.

## Introduction

Thyrotropin (thyroid-stimulating hormone, TSH) is a pivotal activators of the production of thyroid hormones (T3 and its prohormone, T4) from the thyroid gland. TSH is a heterodimer consisting of an  $\alpha$  chain (chorionic gonadotropin  $\alpha$ , CGA) and a  $\beta$  chain (TSH $\beta$ ) [1]. TSH $\beta$  determines the hormonal specificity of TSH whereas CGA heterodimerizes with CG $\beta$ , follicle stimulating hormone (FSH)  $\beta$  and luteinizing hormone (LH)  $\beta$  subunits, to form CG, FSH and LH, respectively [2]. Two transcription factors, GATA2 and Pit1, are the critical determinants of thyrotroph differentiation in pituitary [3] and they directly bind with the DNA sequence of the TSH $\beta$  promoter [2, 4]. Our previous study revealed that the actual activator of the TSH $\beta$  gene is GATA2, whereas Pit1 is necessary for protecting the function of GATA2 from the suppressor region (SR), which is located immediately downstream to the GATA-responsive elements (GATA-REs) [5]. Although detailed analysis has not been performed, hypothyroidism in patients with mutant GATA2 has been reported [6, 7]. Previously we have reported that the signal of thyrotropin releasing hormone (TRH) from hypothalamic paraventricular nucleus (PVN) enhanced the transcriptional activity of GATA2 via the protein kinase C (PKC) pathway [8]. We also found that, although this signaling causes the further stimulation of the TSH $\beta$  and CGA expression in the thyrotroph, the T3-dependent repression is dominant over the activation via this pathway.

The mechanisms underlying the T3 receptor (TR)-mediated transcriptional activation of T3-target genes (positive regulation) have been well characterized [9]. In these studies, monkey kidney-derived CV1 cells [10] have often been utilized [11–13]. In contrast, the mechanism underlying the T3-dependent transcriptional repression via TR (negative regulation) has not yet been clarified. TSH $\beta$  and CGA genes are the typical genes, which are negatively regulated by T3 [14]. In the mice deficient in TR $\beta$ 2 [15], which is the pituitary-specific TR, T3-dependent inhibition is ameliorated. Thus, TR $\beta$ 2 is considered to be the major TR subtype mediating the negative regulation of the TSH $\beta$  and CGA genes. As an analogy of the T3-responsive element (TRE) in the positively regulated genes, the presence of the so-called negative TRE (nTRE) in the TSH $\beta$  gene has long been postulated [1, 16]. Previously we reported that T3-dependent inhibition of the TSH $\beta$  promoter is readily detected even in CV1 cells as long as GATA2, Pit1 and TRs are co-expressed [17]. Unexpectedly, this system revealed that inhibition of the TSH $\beta$  gene by T3 is maintained even after the deletion or mutation of the reported nTRE [18]. Thus, instead of the nTRE model, we proposed the tethering model in which TR $\beta$ 2 interacts with GATA2 Zn-fingers via protein-protein interaction and interferes with the transcriptional function of GATA2 in a T3-dependent manner [13, 18], as observed in the case of inhibition of NF $\kappa$ B-induced transactivation by liganded glucocorticoid receptor [19]. The CGA promoter is also stimulated by GATA2 via its GATA-RE and inhibited by liganded TR $\beta$ 2 [8, 18]. Thus, in the transcriptional regulation of the TSH $\beta$  and CGA genes, GATA2 functions as the platform to select either the activating signal by TRH or inhibitory signal by liganded TR $\beta$ 2 [8, 13].


Importantly, the serum TSH concentrations are drastically reduced by a small increase of thyroid hormone levels [20]. In the clinical studies, values of serum TSH and thyroid

hormones are usually plotted in a log scale and linear scale, respectively [21–24]. Thus, this feature, often referred to as a “log-linear relationship”, makes the serum TSH concentration a very sensitive indicator of thyroid gland function [25]. Although numerous mathematical models [26–28] and the involvement of genetic factors [29] have been proposed, there are few experimental systems that enable us to explore the molecular mechanism underlying this unique relationship between them. GATA2 is expressed not only in thyrotroph but also in gonadotroph [2, 3, 30] and placenta [31–33], resulting in the potentiation of the CGA gene expression [34, 35]. Using trophoblast stem cell line, Rcho-1, and TS cells, Ray et al. [36] surveyed approximately 100-kb regions of the rat and mouse GATA loci using quantitative chromatin immunoprecipitation (ChIP) assay. They demonstrated that the GATA2 gene is transactivated by its own translation product (i.e., GATA2 protein) via multiple GATA-REs in this gene (Fig 1A), as in the case of hematopoietic cell lineage [37–39]. Hence, GATA2 expression is controlled by intracellular positive feedback autoregulation [39]. This mechanism of the GATA2 gene also appears to play a role in pituitary because the thyrotroph and gonadotroph-specific expression of dominant negative-type GATA2 mutant down-regulates the expression of endogenous GATA2 *in vivo* [3]. Given that intracellular positive feedback autoregulation of the GATA2 gene is non-linear, we speculated that inhibition of this control system by T3 may display an inverse non-linear pattern as observed in clinical cases. The GATA2 gene is driven by two promoter, 1S and 1G (Fig 1) [40, 41]. Both promoters are activated by GATA2 via the aforementioned GATA-REs [42] and their transcripts can be detected in the gonadotroph cell line, L $\beta$ T2 [30, 43, 44]. Here, we constructed the chloramphenicol acetyltransferase (CAT)-reporter genes, in which the 1S-promoter was fused to reported GATA-REs derived from the GATA2 gene [36, 39, 45], and co-transfected them with expression plasmids for GATA2 and TR $\beta$ 2 into CV1 cells. We found that liganded TR $\beta$ 2 represses the GATA2-induced transcriptional function via these GATA-REs. Multi-dimensional input function theory suggested that liganded TR $\beta$ 2 functions as a classical transcriptional repressor of the GATA2 promoter. Next, we evaluated the effect of T3 on the endogenous GATA2 protein expression in L $\beta$ T2 cells. We found that T3 reduces GATA2 protein expression independently of the ubiquitin-proteasome system. We observed robust reduction in GATA2 mRNA expression by T3 at concentrations corresponding to those between mild hypothyroidism and euthyroidism. Thus, this study may provide the important insights for the unique relationship between serum TSH and thyroid hormones *in vivo*.

## Materials and methods

### Plasmid constructions

Expression plasmids for rat TR $\beta$ 2 (pCMX-rTR $\beta$ 2) and mouse GATA2 (pcDNA3-mGATA2) have been described previously [8, 17, 18]. Because the firefly luciferase-based reporter gene may be artificially suppressed by liganded TR [[13, 46] and references therein], we employed a CAT-based reporter gene. Three luciferase-based reporter plasmids (generous gifts from Drs. Emery H. Bresnick and Meghan E. Boyer), 1S-Luc (GATA2 gene-derived 1S promoter), (-3.9) 1S-Luc (-3.9 kb GATA-RE fused with 1S-promoter) and (+9.5)1S-Luc (+9.5 kb GATA-RE fused with 1S-promoter) were amplified by polymerase chain reaction (PCR) using three forward primers, 1Spro-UE2 (5' -gggggaattcgccagaaagcccctgtctggggac-3'), 3.9-UME2 (5' -gggggaattcacgcgaagccgccaggtg-3') or 9.5-UME2: (5' -gggggaattcacgcgtccccgcagctaccgggacccccctcctct-3') and a reverse primer, 1Spro-DNB2 (5' -ggggcatatgagatctgggagacctgagcagtgag-3'). These PCR products were digested by EcoRI and Bgl-II. The human D2 promoter in hD2-CAT [47] was replaced by these PCR products after digestion with restriction enzymes (EcoRI and Bgl-II),



**Fig 1. Structures of the mouse GATA2 gene and CAT-reporter genes.** (A) In the mouse GATA2 gene, there are at least four GATA-REs as well as two promoters (1S and 1G) that are regulated by these GATA-REs. 1S-promoter may also contain a functional GATA-RE (see main text). (B) 1S promoter, promoter fused with -3.9 GATA-RE or promoter fused with +9.5 GATA-RE were subcloned to the CAT reporter gene to generate 1S-CAT, (-3.9)1S-CAT and (+9.5)1S-CAT, respectively.

<https://doi.org/10.1371/journal.pone.0227646.g001>

generating 1S-CAT, (-3.9)1S-CAT and (+9.5)1S-CAT, respectively (Fig 1B). In these constructs, the pUC-derived AP-1-like sequence was deleted because it might also mediate artificial T3-dependent inhibition [[13, 46] and references therein]. All subcloning sites were confirmed by sequencing.

### Cell culture and transient transfection

CV1 cells [10] were grown in a monolayer culture at 37°C under CO<sub>2</sub>/air (1:19) in Dulbecco's modified Eagle's medium (DMEM) containing 10% (v/v) fetal calf serum (FCS), penicillin G (100 units/ml), and streptomycin (100 µg/ml). CV1 cells were trypsinized and plated in six-well plates for 24 hr prior to transient transfection using the calcium-phosphate technique [17]. Cells at a density of 2 × 10<sup>5</sup> cells per well were transfected with 1.0 µg of the CAT reporter genes (see above), pcDNA3-mGATA2 and pCMX-rTRβ2 together with 0.9 µg of the β-galactosidase expression vector, pCH111 (a modified version of pCH110, Pharmacia LKB Biotechnology, Piscataway, NJ, USA). The total amount of expression plasmid was adjusted with the empty pCMX vector (3.6 µg of DNA in total per dish). After exposure to calcium phosphate/DNA precipitates for 20 hr, the medium was replaced with fresh DMEM containing 10% FCS depleted of thyroid hormones [[17] and references therein] or medium supplemented with T3.

Cells were harvested after incubation for an additional 24 hr, and CAT activity was measured as described previously [17]. Mouse thyrotroph-derived T $\alpha$ T1 cells [48] (a kind gift from Dr. Pamela Mellon, University of California, CA, USA) were seeded on Matrigel-coated plates (Becton Dickinson Labware, Bedford, MA, USA). L $\beta$ T2 cells, a mouse gonadotroph cell line [43], were cultured in DMEM supplemented with 10% FCS. T $\alpha$ T1 cells and L $\beta$ T2 cells were maintained under the same conditions as for CV1 cells [17].

### Theoretical characterization of liganded TR $\beta$ 2 as transcriptional repressor

Based on the theory of multi-dimensional input function (Fig 3, inset) proposed by Alon [49], we calculated the data from CAT reporter assay (Fig 3) in CV1 cells transfected with fixed amount of (+9.5)1S-CAT and pCMX-rTR $\beta$ 2 together with various amount of pcDNA3-m-GATA2 (0 to 0.2  $\mu$ g/dish) in the presence of 0–1000 nM T3, using Mathematica software (Champaign, IL, USA). The concentration of liganded TR $\beta$ 2 was represented as the T3 concentration ([T3]).

### Western blot analysis


Same amounts (200  $\mu$ l/dish) of whole cell extracts of T $\alpha$ T1 cells or L $\beta$ T2 cells cultured in a 10 cm dish in the presence or absence of 10  $\mu$ M MG132 (ChemScene, Monmouth Junction, NJ, USA) and/or T3 for 24 hr were fractionated (20  $\mu$ l/lane) by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), and then, subjected to Western blot analysis with an anti-GATA2 monoclonal antibody (a kind gift from Drs. Yasuharu Kanki and Tatsuhiko Kodama (University of Tokyo and Perseus Proteomics Inc, Tokyo, Japan)). To assess the levels of transfected GATA2, CV1 cells in a 6-cm dish were transfected with pcDNA3-mGATA2 (5  $\mu$ g/dish) using calcium-phosphate technique. After incubation for an additional 24 hr, the cells were harvested and subjected to Western blot analysis as mentioned above.

### Reverse transcription-quantitative polymerase chain reaction (RT-qPCR)

L $\beta$ T2 cells cultured in 10% FCS were incubated with various concentration of T3, and total RNAs was purified using the acid guanidinium thiocyanate-phenol-chloroform extraction method [50]. Total RNA ( $\mu$ g) was mixed with random hexanucleotides and 200 units of Moloney murine leukemia virus reverse transcriptase (Invitrogen Corp., Carlsbad, CA, USA) for first-strand cDNA synthesis. Using the SYBR Green I kit and a LightCycler (Roche Diagnostics, Mannheim, Germany), precipitated cDNA was quantified by real-time PCR using the following primers: forward primer for exon 1G (5' – CACCCCTATCCCGTGAATCCG–3') and reverse primer for exon 2 (5' – AGCTGTGCTGCCTCCATGTAGTTAT–3') [40]. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was also amplified from cDNA using a forward primer (5' – TGAACGGAAGCTCACTGG–3') and reverse primer (5' – TCCACCACCCTGT TGGCTGTA–3'). The thermal cycling conditions were as follows: 10 min at 95°C, followed by 50 cycles of 10 s at 95°C for denaturing, 10 s at 62°C for annealing, and 7 s at 72°C for extension [50]. PCR signals were quantitatively analyzed using LightCycler software version 3.5 (Roche Diagnostics, Basel, Switzerland).

### Statistical analysis

The CAT reporter assay with CV1 cells and RT-qPCR with L $\beta$ T2 cells were performed in duplicate three or more times, and each result was expressed as the mean  $\pm$  S.E. Statistical significance was examined by ANOVA and Fisher's protected least significant difference test




**Fig 3. The liganded TRβ2 functions as a classical transcriptional repressor defined according to multi-dimensional input function theory.** (+9.5)1S-CAT was transfected into CV1 cells along with the expression plasmid for TRβ2 (0.2 μg) and various amounts of that for GATA2 (0 to 200 ng/dish) and treated with various concentrations of T3 (0 to 1000 nM). The data are plotted as a 3-dimensional graph. Results indicate that the liganded TRβ2 tethered to GATA2 Zn-finger functions as a classical transcriptional repressor (inset). The concentration of T3 ([T3]) represents that of liganded TRβ2. Using the data in (A) and the formula for multi-dimensional input function theory (inset) proposed by Alon [49], we calculated the DNA association affinity of GATA2 (K1), liganded TRβ2 (represented as [T3], K2), production rate of the promoter (βz) and basal transcriptional activity (leakage) of the promoter (β'z) as 0.809, 0.081, 544.33 and 129.32, respectively. G, amount of the expression plasmid for GATA2.

<https://doi.org/10.1371/journal.pone.0227646.g003>

using Stat View 4.0 software (Abacus Concepts, Berkeley, CA, USA). A P value <0.05 was considered statistically significant.

### Results

According to the ChIP analysis of the 100-kb regions of the rat and mouse GATA2 loci [36–38], the GATA2 gene harbors multiple GATA-REs (Fig 1A). In trophoblasts [36] and hematopoietic cells [39], GATA2 protein on these DNA elements activates the transcription of this gene from 1S and 1G promoters, resulting in positive feedback autoregulation [39]. Among these GATA-REs, the functions of -3.9 kb and +9.5 kb GATA-REs have been intensively characterized [36, 39, 51]. The expression plasmids of GATA2 and TRβ2 were co-transfected with the CAT reporter genes, of which 1S-promoter was fused to these GATA-REs (Fig 1B) into CV1 cells, which lack endogenous GATA2 [18] or TRs [12]. As shown in Fig 2A, GATA2




**Fig 2. In CV1 cells, liganded TRβ2 negatively regulates the GATA2-promoter induced by GATA2 itself.** (A) Using the calcium phosphate method, 2.0 μg of 1S-CAT, (-3.9)1S-CAT or (+9.5)1S-CAT was transfected into CV1 cells that were plated at a density of 2×10<sup>5</sup> cells per well in a six-well plate along with the expression plasmids for TRβ2 (0.2 μg) and GATA2 (0.1 μg) in the presence or absence of 100 nM T3. \*, P<0.05 and \*\*, P<0.01 for vector vs. GATA2 expression plasmids. #, P<0.05 for vehicle vs. 100 nM T3. CAT activity for pCMV-CAT (5.0 ng/well) was taken as 100%. Data are expressed as the mean ± S.E. of three to five independent experiments. (B) (+9.5) 1S-CAT was transfected into CV1 cells as shown in (A) and the cells were treated with various concentration of T3. The results are plotted with [T3] (x-axis) in a log scale and the CAT activity (y-axis) in a linear scale. (C) The data same in (B) are plotted with both [T3] and the CAT activity plotted in a linear scale. \*, P<0.05 and \*\*, P<0.01 vs. T3 (-).

<https://doi.org/10.1371/journal.pone.0227646.g002>

significantly activated 1S-CAT and (+9.5)1S-CAT. The GATA2-dependent activation of 1S-CAT is in agreement with a previous report showing that 1S-promoter also harbors several putative GATA-REs [40]. As shown in Fig 2A, T3 significantly repressed the activities of 1S-CAT and (+9.5)1S-CAT in the presence of GATA2 and TRβ2 (P<0.05). (-3.9)1S-CAT also exhibited a tendency to be activated by GATA2 and suppressed by liganded TRβ2 although the levels of reduction were not statistically significant. As shown in Fig 2B, the transcriptional activity of (+9.5)1S-CAT decreased to approximately 40% by T3 addition. Although Fig 2B was plotted with the T3 concentration ([T3]) in the log scale and CAT activity in the linear scale, drastic repression by T3 was readily observed when both axes were plotted in the linear scale (Fig 2C). Using the experimental system in Fig 2B, we aimed to clarify whether the liganded TR functions as a transcriptional repressor that was defined by the theory of multi-





**Fig 4. Endogenous protein expression of GATA2 is inhibited by T3 and enhanced by TPA in LβT2 cells.** (A) Western blot with anti-GATA2 antibody revealed that expression of endogenous GATA2 protein in LβT2 cells is higher than that in TαT1 cells. The GATA2 protein in both cell lines was reduced after 24 hr incubation with 100 nM T3. (B) Western blot with anti-TRβ2 antibody for the whole cell extract of LβT2 cells or that of CV1 cells transfected with expression plasmid of rat TRβ2. (C) FLAG-tagged GATA2 gene driven by the CMV-promoter was transfected into LβT2 cells in 10-cm dish and Western blot with anti-FLAG antibody was performed with the whole cell extract. (D) Western blot with anti-GATA2 antibody for the whole cell extract of LβT2 cells plated in a 10-cm dish treated with 0 to 10 nM TPA for 24 hr.

<https://doi.org/10.1371/journal.pone.0227646.g004>

dimensional input function proposed by Uri Alon (Fig 3 in set) [49]. We co-transfected various amounts of expression plasmids for GATA2 with a fixed amount of (+9.5)IS-CAT into CV1 cells, and evaluated the effect of T3 using 3-dimensional graph (Fig 3). In this setting, we postulated that [T3] represents the concentration of liganded TRβ2. We were able to calculate the promoter association affinity of GATA2 (K1), liganded TRβ2 (K2) represented by [T3], production rate of the promoter (βz) and basal transcriptional activity (leakage) of the promoter (β'z) as 0.809, 0.081, 544.33 and 129.32, respectively. This finding indicated that the liganded TRβ2 tethered to GATA2 behaves as a classical transcriptional repressor.

Next, we examined the inhibitory effect of liganded TRβ2 in the cells, where the GATA2 gene is endogenously expressed presumably via its positive feedback loop. While TαT1 is known to be a thyrotroph cell line [48], its GATA2 protein expression level is modest [8]; we found that the suppressive effect of T3 on TSHβ and the CGA mRNA was mild (data not shown) and completely lost, respectively. Other candidates are gonadotroph-derived LβT2 [43] and choriocarcinoma-derived JEG3 [52], both of which express the CGA gene [30, 34, 35, 52]. Because the expression level of TR in JEG3 cells is very low (personal communication from Dr. Takashi Nagaya, Nagoya University in Japan), we compared expression of the endogenous GATA2 protein in LβT2 cells [53] with that in TαT1 cells by Western blot analysis using anti-GATA2 antibody. As shown in Fig 4A, we observed higher GATA2 protein expression in LβT2 cells than that in TαT1 cells. Both bands were repressed by treatment with 100 nM T3, suggesting the presence of TRs. As expected, we observed endogenous TRβ2 in LβT2 cells (Fig


4B) as observed in T $\alpha$ T1 cells [17]. In contrast, 100 nM T3 did not affect the level of CMV-promoter driven FLAG-tagged GATA2, which was transfected into L $\beta$ T2 cells and was detected using anti-FLAG antibody (Fig 4C). Based on these findings, we employed L $\beta$ T2 cells in the subsequent experiments. As is the case of PKC-dependent potentiation of GATA2-induced transactivation in the TSH $\beta$ , CGA [8] and vascular adhesion molecule-1 genes [54], 12-O-tetradecanoylphorbol-13-acetate (TPA), a PKC activating compound, enhanced the expression of the GATA2 protein (Fig 4D). Fig 5A and 5B exhibit the T3-dependency and the time course of the GATA2 protein reduction in L $\beta$ T2 cells. The endogenous GATA2 protein was clearly down regulated after 24 hr treatment with 100 nM T3 (Fig 5B). As GATA2 is known to be rapidly degraded by the ubiquitin-proteasome system [55, 56], we examined the GATA2 protein levels in the presence of MG132, a cell-permeable proteasome inhibitor. While longer exposure of MG132 increased the yield of the GATA2 protein (Fig 5C), T3 decreased it even in the presence of this compound (Fig 5D), suggesting that T3-dependent repression of GATA2 protein occurs prior to degradation by the ubiquitin-proteasome system.

Measurement of the GATA2 mRNA levels enables us to directly evaluate the T3-dependent repression of GATA2 gene transcription without the influence of the ubiquitin-proteasome system. Thus, we performed RT-qPCR to evaluate the endogenous expression of the GATA2 mRNA in L $\beta$ T2 cells in the presence of various concentration of T3. When GATA2 mRNA standardized by GAPDH mRNA (GATA2/GAPDH ratio) and T3 concentration was plotted in a linear scale and log scale, respectively (Fig 6A), a significant inhibition was observed; the shape of this relationship was a negative sigmoidal curve, suggesting that T3-dependent repression of the GATA2 gene is non-linear. When both values were plotted in a linear scales, robust reduction in the GATA2 mRNA level by T3 was observed (Fig 6B). Moreover, a rigorous decrease in the GATA2/GAPDH ratio by T3 was still observed when the former was plotted in a log scale and the latter was plotted in a linear scale between 0.1–1.0 nM, the concentrations that correspond to those in moderate hypothyroidism and euthyroidism [57],

## Discussion

GATA2 is a critical transcription factor essential for the gene expression of TSH $\beta$  and CGA gene [3–5]. As illustrated in Fig 7, we demonstrated here that liganded TR $\beta$ 2 not only directly inhibits the promoters of the TSH $\beta$  and CGA genes [17, 18] but also interferes with the GATA2 promoter. In agreement with our findings, two independent studies reported that GATA2 mRNA expression was increased in the pituitary gland of hypothyroid mice [58, 59]. As shown in Fig 3, the liganded TR $\beta$ 2 appears to function as a classical transcriptional repressor [49]. However, liganded TR $\beta$ 2 may cause drastic suppression of the GATA2 gene expression by means of the global interference with the positive feedback loop of this gene (Fig 7), resulting in the maintenance of the homeostasis in hypothalamus-pituitary-thyroid (H-P-T) axis. In mice, deletion of the GATA2 gene specifically in anterior pituitary is not lethal [60] because loss of GATA2 can be partially compensated by the increased expression of the gene for GATA3, another member of the GATA transcription factor family [61]. Interestingly, GATA3 expression may also be controlled via its positive feedback autoregulation [62, 63]. Liganded TRs may also interfere with this GATA3 autoregulation in these mice because they repress GATA3-dependent transactivation [8, 18].

The nTRE was originally defined based on the hypothesis that an unliganded TR may function as transcriptional activator for a gene that is negatively regulated by T3 [16]. However, this hypothesis has been denied because TR-null mice displayed the increased gene expression of the TSH $\beta$  or CGA [15, 64–66]. Furthermore, there are several experimental and theoretical issues in the original report of nTREs (reviewed in [13]). Instead, we proposed the tethering




**Fig 5. T3-dependent repression of GATA2 protein occurs independently of the ubiquitin-proteasome system.** (A) Western blot with anti-GATA2 antibody for the whole cell extract of LβT2 cells plated in a 10-cm dish and treated with 0 to 100 nM T3 for 24 hr. CV1/GATA2: CV1 cells transfected with expression plasmid for mouse GATA2. (B) Western blot with anti-GATA2 antibody for the whole cell extract of LβT2 cells plated in a 10-cm dish treated with 0 to 100 nM T3 for 0 to 24 hr. (C) Western blot with anti-GATA2 antibody for the whole cell extract of LβT2 cells plated in a 10-cm dish and treated with 10 μM MG132 for 0 to 6 hr before harvest. (D) Western blot with anti-GATA2 antibody for the whole cell extract (15 μL/lane or 30 μL/lane) of LβT2 cells plated in a 10-cm dish and treated for 24 hr with 10 μM MG132 and 23.5 hr with 100 nM T3 before harvest.

<https://doi.org/10.1371/journal.pone.0227646.g005>

model in which TR associates directly with GATA2 via protein-protein interaction and interferes with GATA2-dependent transactivation in a T3-dependent manner [13, 18]. Although T3 has been known to repress type-2 deiodinase (D2) gene/activity in several organs and cultured cells including thyrotroph [67], no DNA sequence similar to the nTRE reported in the TSHβ gene has been identified in this gene to date [68, 69]. Interestingly, however, the D2 gene harbors the two highly conserved GATA-REs [47]. As expected, our previous work revealed that liganded TRβ2 suppresses the D2 promoter by a tethering mechanism via GATA2 on these GATA-REs [47]. In the GATA2 gene, the critical function of +9.5 kb GATA-RE (Fig 1A) was evidenced based on hematopoietic stem/progenitor cell depletion observed in the mice harboring the deletion of this GATA-RE [45].

Gonadotroph-derived LβT2 cells express the endogenous GATA2 gene (Figs 4 and 5). Thus, these cells may represent a more suitable model than CV1 cells to observe the T3-dependent inhibition of autoregulation of the GATA2 gene. As expected, the graph of T3 inhibition of the GATA2 gene evaluated by RT-qPCR analysis in LβT2 cells is a negative sigmoidal curve (Fig 6A). This pattern of regulation is considered to be suitable for rapid “on and off” control of expression by T3. Fig 6A shows the GATA2/GAPDH ratio (y-axis) in a linear scale and T3 (x-axis) in a log scale, this is the reverse of that seen in clinical situation, i.e., TSH in a log scale and thyroid hormones in a linear scale [21–23, 24]. However, strong repression of GATA2/




**Fig 6. Inhibition of GATA2 mRNA by T3 in LβT2 cells.** (A) Using RT-qPCR with primers specific for exon 1G and exon 2, the effect of T3 on the expression of GATA2 mRNA in LβT2 cells was evaluated. The shape of this graph was a negative sigmoidal curve. (B) The same data as in (A) are plotted with both [T3] (x-axis) and the GATA2/GADPH ratio (y-axis) plotted in a linear scale. \*,  $P < 0.01$  vs. T3(-). #,  $P < 0.01$  vs. T3 = 0.1 nM. (C) The same data as in (A) and (B) with the GATA2/GADPH ratio (y-axis) in a log scale and [T3] (x-axis) between 0–1.0 nM in a linear scale, which corresponds to moderate hypothyroidism and euthyroidism [57].

<https://doi.org/10.1371/journal.pone.0227646.g006>

GADPH ratio by T3 was observed when both are plotted in a linear scale (Fig 6B). Furthermore, potent suppression was found even when the GATA2/GADPH ratio and T3 were plotted in a log scale and linear scale, respectively (Fig 6C), especially at the T3 concentration those observed in moderate hypothyroidism and euthyroidism [57]. Thus, in the LβT2 cells, T3-dependent inhibition of positive feedback autoregulation in the GATA2 gene may contribute, at least in part, to the non-linear relationship between TSH production and thyroid hormones in vivo.

In this study, use of cell culture system enabled us to exclude the influence of TRH from hypothalamic PVN. However, the T3-dependent negative regulation of TRH production should be considered [70, 71] because, in thyrotroph, TRH receptor (TRH-R) signaling potentiates the transactivation function of GATA2 via the PKC pathway, resulting in the enhanced expression of the TSHβ and CGA genes [8] as well as the GATA2 gene (Fig 4D). Notably, the alteration in the level of TRH mRNA level in PVN after thyroidectomy and subsequent T3 supplementation are very similar to those of TSHβ mRNA in the pituitary [72]. Moreover, the



**Fig 7. A schematic representation of hypothalamus-pituitary-thyroid (H-P-T) axis and transcriptional regulation of the GATA2, TSH $\beta$  and CGA genes by GATA2 and liganded TR $\beta$ 2 in thyrotroph.** To simplify, only one GATA-RE is illustrated in the 5' region of the GATA2 gene. In a TRH-dependent manner, TRH-R stimulates the transcriptional activity of GATA2 via the PKC pathway. In contrast, TR $\beta$ 2 associates with the Zn-finger region of GATA2 via protein-protein interaction, resulting in the T3-dependent interference of GATA2-dependent transactivation (tethering mechanism). Our study suggests that liganded TR $\beta$ 2 inhibits not only the transcription of the TSH $\beta$  and CGA genes but also the positive feedback autoregulation of the GATA2 gene. Thus, there are the intracellular cascades of T3-dependent negative regulation between GATA2 and TSH $\beta$ /CGA. Reduction of GATA2 protein expression by T3 appears to occur at transcriptional level and is independent from the ubiquitin-proteasome system. T3 also causes TSH $\beta$  mRNA to be unstable. D2, type 2 deiodinase (the major deiodinase in thyrotroph). SR, suppressor region (see main text). In addition to GATA2, the CGA promoter is regulated by multiple transcription factors including Msx1, SF-1, Lhx3a, cAMP response element binding protein (CREB) and Ptx1 [35].

<https://doi.org/10.1371/journal.pone.0227646.g007>

GATA2 and TR $\beta$ 2 genes are induced in the cultured neurons that are over-expressed with Sim1, a transcription factor that determines the differentiation of the PVN in the hypothalamus [73, 74]. We recently observed that GATA2 is expressed in the TRH neuron of the rat PVN, and found that liganded TR $\beta$ 2 represses the preproTRH gene activated by GATA2 on its GATA-RE (Kuroda et al. paper in preparation). Thus, T3-dependent inhibition of the positive feedback regulation of the GATA2 gene in both thyrotrophs and the PVN may share common mechanism. Indeed, Fekete et al. [71] previously reported an inverse log-linear relationship between the numbers of preproTRH positive cells in rat PVN as evaluated by in situ hybridization analysis and plasma T3 concentration. Interestingly, detailed analysis of 152261 human subjects revealed that the relationship between TSH in a log scale and free T4 in a linear scale

may not be inverse log-linear but can be described by two overlapping negative sigmoidal curves [24].

Several additional mechanisms are considered to be involved in the negative feedback regulation of the H-P-T axis. First, although we focused on the feedback regulation of the GATA2 gene, T3-dependent inhibition may be amplified by the intracellular cascades between the GATA2 gene and the TSH $\beta$  and CGA genes (Fig 7). Second, T3 also represses the expression of prohormone convertase 1/3 and 2, which play essential roles in the processing of the proTRH peptide in the PVN [75], and the TRH-R expression in pituitary [76, 77]. Third, T3 destabilizes the TSH $\beta$  mRNA but not CGA mRNA [78]. Fourth, T3 may inhibit the expression of Pit1 [79], which is a critical transcription factor for the TSH $\beta$  gene (Fig 7), but not the CGA gene [2]. Finally, the cell cycle of thyrotrophs [80, 81] may also be involved because sustained hypothyroidism often causes thyrotroph hyperplasia, resulting in the massive enlargement of the anterior pituitary [82].

GATA2 was originally identified as an essential transcription factor that mediates differentiation of hematopoietic cells [61]. Consistently, hematopoietic cell differentiation in mice is affected by T3 administration [83]; moreover, it is impaired in the mice that harbor genetic defects for the gene coding TR $\alpha$  [83, 84], the major TR subtype in the erythroid cell lineage. In human subjects with mutations in the TR $\alpha$  gene, anemia and erythroid defect were reported [85]. Thus, the precise mechanism of the effect of T3 on the GATA2-related hematopoiesis should be investigated in future. Current results also predict that serum TSH level may be correlated with placental function because this organ expresses GATA family members [32, 33] as well as TRs [86]. As is the case with the TSH $\beta$  and CGA gene in the pituitary (Fig 7), GATA2 and GATA3 regulate the expression of the genes that encode the critical proteins responsible for maintaining the placental function, including the component of bone morphogenetic protein 4, Nodal and Wnt signals [32]. These findings may provide the insight into why subtle elevation in the serum TSH concentration ( $> 2.5 \mu\text{IU/L}$ ) is correlated with the pregnancy loss rate in thyroid antibody-negative women [87–89].

## Supporting information

### S1 Raw Images.

(PDF)

## Acknowledgments

We are grateful to the following investigators for providing plasmids: Drs. Meghan E. Boyer and Emery H. Bresnick (University of Wisconsin, USA), Kazuhiko Umesono (Kyoto University, Japan) and Masayuki Yamamoto (Tsukuba University, Japan). We also thank Drs. Yasuharu Kanki and Tatsuhiko Kodama (University of Tokyo) for providing the anti-GATA2 antibody and Dr. Pamela Mellon (University of California, CA, USA) for the T $\alpha$ T1 and L $\beta$ T2 cells.

## Author Contributions

**Conceptualization:** Shigekazu Sasaki.

**Data curation:** Hiroko Misawa Nakamura, Shigekazu Sasaki.

**Formal analysis:** Hiroko Misawa Nakamura, Shigekazu Sasaki, Akio Matsushita, Kenji Ohba, Go Kuroda, Hiroshi Haeno, Takuhiro Nishio, Shuichi Yoshida, Yutaka Oki, Takafumi Suda.

**Funding acquisition:** Shigekazu Sasaki.

**Investigation:** Naoko Hirahara, Hiroko Misawa Nakamura, Shigekazu Sasaki, Go Kuroda, Yuki Sakai, Shinsuke Shinkai.

**Methodology:** Naoko Hirahara, Hiroko Misawa Nakamura, Akio Matsushita, Hiroshi Haeno.

**Project administration:** Takafumi Suda.

**Resources:** Shigekazu Sasaki, Akio Matsushita, Go Kuroda, Yuki Sakai, Shinsuke Shinkai.

**Software:** Takuhiro Nishio, Shuichi Yoshida.

**Supervision:** Shigekazu Sasaki.

**Validation:** Shigekazu Sasaki, Akio Matsushita, Kenji Ohba, Go Kuroda, Yuki Sakai, Shinsuke Shinkai, Hiroshi Haeno, Takuhiro Nishio, Shuichi Yoshida, Yutaka Oki, Takafumi Suda.

**Visualization:** Shigekazu Sasaki.

**Writing – original draft:** Naoko Hirahara, Hiroko Misawa Nakamura, Shigekazu Sasaki.

**Writing – review & editing:** Naoko Hirahara.

## References

1. Cohen RN, Wondisford FE. Chemistry and Biosynthesis of Thyrotropin. Werner and Ingbar's The Thyroid. 2012; Tenth Edition: pp149–61.
2. Rizzoti K. Genetic regulation of murine pituitary development. *J Mol Endocrinol*. 2015; 54(2):R55–73. Epub 2015/01/15. <https://doi.org/10.1530/JME-14-0237> PMID: 25587054
3. Dasen JS, O'Connell SM, Flynn SE, Treier M, Gleiberman AS, Szeto DP, et al. Reciprocal interactions of Pit1 and GATA2 mediate signaling gradient-induced determination of pituitary cell types. *Cell*. 1999; 97(5):587–98. [https://doi.org/10.1016/s0092-8674\(00\)80770-9](https://doi.org/10.1016/s0092-8674(00)80770-9) PMID: 10367888
4. Gordon DF, Lewis SR, Haugen BR, James RA, McDermott MT, Wood WM, et al. Pit-1 and GATA-2 interact and functionally cooperate to activate the thyrotropin beta-subunit promoter. *The Journal of biological chemistry*. 1997; 272(39):24339–47. <https://doi.org/10.1074/jbc.272.39.24339> PMID: 9305891
5. Kashiwabara Y, Sasaki S, Matsushita A, Nagayama K, Ohba K, Iwaki H, et al. Functions of PIT1 in GATA2-dependent transactivation of the thyrotropin beta promoter. *J Mol Endocrinol*. 2009; 42(3):225–37. <https://doi.org/10.1677/JME-08-0099> PMID: 19103719
6. Spinner MA, Sanchez LA, Hsu AP, Shaw PA, Zerbe CS, Calvo KR, et al. GATA2 deficiency: a protean disorder of hematopoiesis, lymphatics, and immunity. *Blood*. 2014; 123(6):809–21. <https://doi.org/10.1182/blood-2013-07-515528> PMID: 24227816
7. Donadieu J, Lamant M, Fieschi C, de Fontbrune FS, Caye A, Ouachee M, et al. Natural history of GATA2 deficiency in a survey of 79 French and Belgian patients. *Haematologica*. 2018; 103(8):1278–87. <https://doi.org/10.3324/haematol.2017.181909> PMID: 29724903
8. Ohba K, Sasaki S, Matsushita A, Iwaki H, Matsunaga H, Suzuki S, et al. GATA2 mediates thyrotropin-releasing hormone-induced transcriptional activation of the thyrotropin beta gene. *PloS one*. 2011; 6(4): e18667. <https://doi.org/10.1371/journal.pone.0018667> PMID: 21533184
9. Cheng SY, Leonard JL, Davis PJ. Molecular aspects of thyroid hormone actions. *Endocrine reviews*. 2010; 31(2):139–70. <https://doi.org/10.1210/er.2009-0007> PMID: 20051527
10. Jensen FC, Girardi AJ, Gilden RV, Koprowski H. Infection of Human and Simian Tissue Cultures with Rous Sarcoma Virus. *Proceedings of the National Academy of Sciences of the United States of America*. 1964; 52:53–9. <https://doi.org/10.1073/pnas.52.1.53> PMID: 14192657
11. Umesonon K, Evans RM. Determinants of target gene specificity for steroid/thyroid hormone receptors. *Cell*. 1989; 57(7):1139–46. [https://doi.org/10.1016/0092-8674\(89\)90051-2](https://doi.org/10.1016/0092-8674(89)90051-2) PMID: 2500251
12. Umesonon K, Murakami KK, Thompson CC, Evans RM. Direct repeats as selective response elements for the thyroid hormone, retinoic acid, and vitamin D3 receptors. *Cell*. 1991; 65(7):1255–66. [https://doi.org/10.1016/0092-8674\(91\)90020-y](https://doi.org/10.1016/0092-8674(91)90020-y) PMID: 1648450
13. Sasaki S, Matsushita A, Kuroda G, Nakamura HM, Oki Y, Suda T. The Mechanism of Negative Transcriptional Regulation by Thyroid Hormone: Lessons From the Thyrotropin beta Subunit Gene. *Vitam Horm*. 2018; 106:97–127. <https://doi.org/10.1016/bs.vh.2017.06.006> PMID: 29407449

14. Shupnik MA, Ridgway EC, Chin WW. Molecular biology of thyrotropin. *Endocrine reviews*. 1989; 10(4):459–75. <https://doi.org/10.1210/edrv-10-4-459> PMID: 2693083
15. Abel ED, Boers ME, Pazos-Moura C, Moura E, Kaulbach H, Zakaria M, et al. Divergent roles for thyroid hormone receptor beta isoforms in the endocrine axis and auditory system. *The Journal of clinical investigation*. 1999; 104(3):291–300. <https://doi.org/10.1172/JCI6397> PMID: 10430610
16. Wondisford FE, Farr EA, Radovick S, Steinfeldler HJ, Moates JM, McClaskey JH, et al. Thyroid hormone inhibition of human thyrotropin beta-subunit gene expression is mediated by a cis-acting element located in the first exon. *The Journal of biological chemistry*. 1989; 264(25):14601–4. PMID: 2768233
17. Nakano K, Matsushita A, Sasaki S, Misawa H, Nishiyama K, Kashiwabara Y, et al. Thyroid-hormone-dependent negative regulation of thyrotropin beta gene by thyroid hormone receptors: study with a new experimental system using CV1 cells. *Biochem J*. 2004; 378(Pt 2):549–57. <https://doi.org/10.1042/BJ20031592> PMID: 14611644
18. Matsushita A, Sasaki S, Kashiwabara Y, Nagayama K, Ohba K, Iwaki H, et al. Essential role of GATA2 in the negative regulation of thyrotropin beta gene by thyroid hormone and its receptors. *Molecular endocrinology (Baltimore, Md)*. 2007; 21(4):865–84.
19. Petta I, Dejager L, Ballegeer M, Lievens S, Tavernier J, De Bosscher K, et al. The Interactome of the Glucocorticoid Receptor and Its Influence on the Actions of Glucocorticoids in Combatting Inflammatory and Infectious Diseases. *Microbiology and molecular biology reviews: MMBR*. 2016; 80(2):495–522. <https://doi.org/10.1128/MMBR.00064-15> PMID: 27169854
20. Reichlin S, Utiger RD. Regulation of the pituitary-thyroid axis in man: relationship of TSH concentration to concentration of free and total thyroxine in plasma. *The Journal of clinical endocrinology and metabolism*. 1967; 27(2):251–5. <https://doi.org/10.1210/jcem-27-2-251> PMID: 4163614
21. Wehmann RE, Nisula BC. Radioimmunoassay of human thyrotropin: analytical and clinical developments. *Critical reviews in clinical laboratory sciences*. 1984; 20(3):243–83. <https://doi.org/10.3109/10408368409165776> PMID: 6373146
22. Spencer CA, LoPresti JS, Patel A, Guttler RB, Eigen A, Shen D, et al. Applications of a new chemiluminometric thyrotropin assay to subnormal measurement. *The Journal of clinical endocrinology and metabolism*. 1990; 70(2):453–60. <https://doi.org/10.1210/jcem-70-2-453> PMID: 2105333
23. Hoermann R, Eckl W, Hoermann C, Larisch R. Complex relationship between free thyroxine and TSH in the regulation of thyroid function. *European journal of endocrinology*. 2010; 162(6):1123–9. <https://doi.org/10.1530/EJE-10-0106> PMID: 20299491
24. Hadlow NC, Rothacker KM, Wardrop R, Brown SJ, Lim EM, Walsh JP. The relationship between TSH and free T(4) in a large population is complex and nonlinear and differs by age and sex. *The Journal of clinical endocrinology and metabolism*. 2013; 98(7):2936–43. <https://doi.org/10.1210/jc.2012-4223> PMID: 23671314
25. Larsen PR, Davies TF, Schlumberger MJ, Hay ID. *Thyroid Physiology and Diagnostic Evaluation of Patients with Thyroid Disorders*. Williams Textbook of Endocrinology. 2015; Thirteenth Edition:pp334–68.
26. Leow MK, Goede SL. The homeostatic set point of the hypothalamus-pituitary-thyroid axis—maximum curvature theory for personalized euthyroid targets. *Theor Biol Med Model*. 2014; 11:35. <https://doi.org/10.1186/1742-4682-11-35> PMID: 25102854
27. Hoermann R, Midgley JEM, Larisch R, Dietrich JWC. Advances in applied homeostatic modelling of the relationship between thyrotropin and free thyroxine. *PloS one*. 2017; 12(11):e0187232. <https://doi.org/10.1371/journal.pone.0187232> PMID: 29155897
28. Dietrich JW, Midgley JEM, Hoermann R. Editorial: "Homeostasis and Allostasis of Thyroid Function". *Front Endocrinol (Lausanne)*. 2018; 9:287.
29. Medici M, Visser WE, Visser TJ, Peeters RP. Genetic determination of the hypothalamic-pituitary-thyroid axis: where do we stand? *Endocrine reviews*. 2015; 36(2):214–44. <https://doi.org/10.1210/er.2014-1081> PMID: 25751422
30. Laverriere JN, L'Hote D, Tabouy L, Schang AL, Querat B, Cohen-Tannoudji J. Epigenetic regulation of alternative promoters and enhancers in progenitor, immature, and mature gonadotrope cell lines. *Molecular and cellular endocrinology*. 2016; 434:250–65. <https://doi.org/10.1016/j.mce.2016.07.010> PMID: 27402603
31. Bai H, Sakurai T, Godkin JD, Imakawa K. Expression and potential role of GATA factors in trophoblast development. *J Reprod Dev*. 2013; 59(1):1–6. <https://doi.org/10.1262/jrd.2012-100> PMID: 23428586
32. Home P, Kumar RP, Ganguly A, Saha B, Milano-Foster J, Bhattacharya B, et al. Genetic redundancy of GATA factors in the extraembryonic trophoblast lineage ensures the progression of preimplantation and postimplantation mammalian development. *Development (Cambridge, England)*. 2017; 144(5):876–88.



33. Paul S, Home P, Bhattacharya B, Ray S. GATA factors: Master regulators of gene expression in trophoblast progenitors. *Placenta*. 2017; 60 Suppl 1:S61–s6.
34. Steger DJ, Hecht JH, Mellon PL. GATA-binding proteins regulate the human gonadotropin alpha-subunit gene in the placenta and pituitary gland. *Molecular and cellular biology*. 1994; 14(8):5592–602. <https://doi.org/10.1128/mcb.14.8.5592> PMID: 7518566
35. Jorgensen JS, Quirk CC, Nilson JH. Multiple and overlapping combinatorial codes orchestrate hormonal responsiveness and dictate cell-specific expression of the genes encoding luteinizing hormone. *Endocrine reviews*. 2004; 25(4):521–42. <https://doi.org/10.1210/er.2003-0029> PMID: 15294880
36. Ray S, Dutta D, Rumi MA, Kent LN, Soares MJ, Paul S. Context-dependent function of regulatory elements and a switch in chromatin occupancy between GATA3 and GATA2 regulate Gata2 transcription during trophoblast differentiation. *The Journal of biological chemistry*. 2009; 284(8):4978–88. <https://doi.org/10.1074/jbc.M807329200> PMID: 19106099
37. Kobayashi-Osaki M, Ohneda O, Suzuki N, Minegishi N, Yokomizo T, Takahashi S, et al. GATA motifs regulate early hematopoietic lineage-specific expression of the Gata2 gene. *Molecular and cellular biology*. 2005; 25(16):7005–20. <https://doi.org/10.1128/MCB.25.16.7005-7020.2005> PMID: 16055713
38. Grass JA, Jing H, Kim SI, Martowicz ML, Pal S, Blobel GA, et al. Distinct functions of dispersed GATA factor complexes at an endogenous gene locus. *Molecular and cellular biology*. 2006; 26(19):7056–67. <https://doi.org/10.1128/MCB.01033-06> PMID: 16980610
39. Bresnick EH, Katsumura KR, Lee HY, Johnson KD, Perkins AS. Master regulatory GATA transcription factors: mechanistic principles and emerging links to hematologic malignancies. *Nucleic Acids Res*. 2012; 40(13):5819–31. <https://doi.org/10.1093/nar/gks281> PMID: 22492510
40. Minegishi N, Ohta J, Suwabe N, Nakauchi H, Ishihara H, Hayashi N, et al. Alternative promoters regulate transcription of the mouse GATA-2 gene. *The Journal of biological chemistry*. 1998; 273(6):3625–34. <https://doi.org/10.1074/jbc.273.6.3625> PMID: 9452491
41. Minegishi N, Ohta J, Yamagiwa H, Suzuki N, Kawauchi S, Zhou Y, et al. The mouse GATA-2 gene is expressed in the para-aortic splanchnopleura and aorta-gonads and mesonephros region. *Blood*. 1999; 93(12):4196–207. PMID: 10361117
42. Grass JA, Boyer ME, Pal S, Wu J, Weiss MJ, Bresnick EH. GATA-1-dependent transcriptional repression of GATA-2 via disruption of positive autoregulation and domain-wide chromatin remodeling. *Proceedings of the National Academy of Sciences of the United States of America*. 2003; 100(15):8811–6. <https://doi.org/10.1073/pnas.1432147100> PMID: 12857954
43. Turgeon JL, Kimura Y, Waring DW, Mellon PL. Steroid and pulsatile gonadotropin-releasing hormone (GnRH) regulation of luteinizing hormone and GnRH receptor in a novel gonadotrope cell line. *Molecular endocrinology (Baltimore, Md)*. 1996; 10(4):439–50.
44. Lo A, Zheng W, Gong Y, Crochet JR, Halvorson LM. GATA transcription factors regulate LHbeta gene expression. *J Mol Endocrinol*. 2011; 47(1):45–58. <https://doi.org/10.1530/JME-10-0137> PMID: 21571865
45. Johnson KD, Hsu AP, Ryu MJ, Wang J, Gao X, Boyer ME, et al. Cis-element mutated in GATA2-dependent immunodeficiency governs hematopoiesis and vascular integrity. *The Journal of clinical investigation*. 2012; 122(10):3692–704. <https://doi.org/10.1172/JCI61623> PMID: 22996659
46. Misawa H, Sasaki S, Matsushita A, Ohba K, Iwaki H, Matsunaga H, et al. Liganded thyroid hormone receptor inhibits phorbol 12-O-tetradecanoate-13-acetate-induced enhancer activity via firefly luciferase cDNA. *PloS one*. 2012; 7(1):e28916. <https://doi.org/10.1371/journal.pone.0028916> PMID: 22253701
47. Matsunaga H, Sasaki S, Suzuki S, Matsushita A, Nakamura H, Nakamura HM, et al. Essential Role of GATA2 in the Negative Regulation of Type 2 Deiodinase Gene by Liganded Thyroid Hormone Receptor beta2 in Thyrotroph. *PloS one*. 2015; 10(11):e0142400. <https://doi.org/10.1371/journal.pone.0142400> PMID: 26571013
48. Yusta B, Alarid ET, Gordon DF, Ridgway EC, Mellon PL. The thyrotropin beta-subunit gene is repressed by thyroid hormone in a novel thyrotrope cell line, mouse T alphaT1 cells. *Endocrinology*. 1998; 139(11):4476–82. <https://doi.org/10.1210/endo.139.11.6283> PMID: 9794455
49. Alon U. *Multi-Dimensional Input-Functions. An Introduction to Systems Biology: Design Principles of Biological Circuits*. 2006; Tenth Edition:pp253–5.
50. Iwaki H, Sasaki S, Matsushita A, Ohba K, Matsunaga H, Misawa H, et al. Essential role of TEA domain transcription factors in the negative regulation of the MYH 7 gene by thyroid hormone and its receptors. *PloS one*. 2014; 9(4):e88610. <https://doi.org/10.1371/journal.pone.0088610> PMID: 24781449
51. Wozniak RJ, Boyer ME, Grass JA, Lee Y, Bresnick EH. Context-dependent GATA factor function: combinatorial requirements for transcriptional control in hematopoietic and endothelial cells. *The Journal of biological chemistry*. 2007; 282(19):14665–74. <https://doi.org/10.1074/jbc.M700792200> PMID: 17347142

52. Kohler PO, Bridson WE. Isolation of hormone-producing clonal lines of human choriocarcinoma. *The Journal of clinical endocrinology and metabolism*. 1971; 32(5):683–7. <https://doi.org/10.1210/jcem-32-5-683> PMID: 5103722
53. Fowkes RC, King P, Burrin JM. Regulation of human glycoprotein hormone alpha-subunit gene transcription in LbetaT2 gonadotropes by protein kinase C and extracellular signal-regulated kinase 1/2. *Biology of reproduction*. 2002; 67(3):725–34. <https://doi.org/10.1095/biolreprod67.3.725> PMID: 12193378
54. Minami T, Abid MR, Zhang J, King G, Kodama T, Aird WC. Thrombin stimulation of vascular adhesion molecule-1 in endothelial cells is mediated by protein kinase C (PKC)-delta-NF-kappa B and PKC-zeta-GATA signaling pathways. *The Journal of biological chemistry*. 2003; 278(9):6976–84. <https://doi.org/10.1074/jbc.M208974200> PMID: 12493764
55. Minegishi N, Suzuki N, Kawatani Y, Shimizu R, Yamamoto M. Rapid turnover of GATA-2 via ubiquitin-proteasome protein degradation pathway. *Genes Cells*. 2005; 10(7):693–704. <https://doi.org/10.1111/j.1365-2443.2005.00864.x> PMID: 15966900
56. Nakajima T, Kitagawa K, Ohhata T, Sakai S, Uchida C, Shibata K, et al. Regulation of GATA-binding protein 2 levels via ubiquitin-dependent degradation by Fbw7: involvement of cyclin B-cyclin-dependent kinase 1-mediated phosphorylation of THR176 in GATA-binding protein 2. *The Journal of biological chemistry*. 2015; 290(16):10368–81. <https://doi.org/10.1074/jbc.M114.613018> PMID: 25670854
57. Ashizawa K, Cheng SY. Regulation of thyroid hormone receptor-mediated transcription by a cytosol protein. *Proceedings of the National Academy of Sciences of the United States of America*. 1992; 89(19):9277–81. <https://doi.org/10.1073/pnas.89.19.9277> PMID: 1409635
58. Costa-e-Sousa RH, Astapova I, Ye F, Wondisford FE, Hollenberg AN. The thyroid axis is regulated by NCoR1 via its actions in the pituitary. *Endocrinology*. 2012; 153(10):5049–57. <https://doi.org/10.1210/en.2012-1504> PMID: 22878400
59. Gergics P, Christian HC, Choo MS, Ajmal A, Camper SA. Gene Expression in Mouse Thyrotrope Adenoma: Transcription Elongation Factor Stimulates Proliferation. *Endocrinology*. 2016; 157(9):3631–46. <https://doi.org/10.1210/en.2016-1183> PMID: 27580811
60. Charles MA, Saunders TL, Wood WM, Owens K, Parlow AF, Camper SA, et al. Pituitary-specific Gata2 knockout: effects on gonadotrope and thyrotrope function. *Molecular endocrinology (Baltimore, Md)*. 2006; 20(6):1366–77.
61. Katsumura KR, Bresnick EH. The GATA factor revolution in hematology. *Blood*. 2017; 129(15):2092–102. Epub 2017/02/10. <https://doi.org/10.1182/blood-2016-09-687871> PMID: 28179282
62. Ranganath S, Murphy KM. Structure and specificity of GATA proteins in Th2 development. *Molecular and cellular biology*. 2001; 21(8):2716–25. <https://doi.org/10.1128/MCB.21.8.2716-2725.2001> PMID: 11283251
63. Nakata Y, Brignier AC, Jin S, Shen Y, Rudnick SI, Sugita M, et al. c-Myb, Menin, GATA-3, and MLL form a dynamic transcription complex that plays a pivotal role in human T helper type 2 cell development. *Blood*. 2010; 116(8):1280–90. <https://doi.org/10.1182/blood-2009-05-223255> PMID: 20484083
64. Forrest D, Hanebuth E, Smeyne RJ, Everds N, Stewart CL, Wehner JM, et al. Recessive resistance to thyroid hormone in mice lacking thyroid hormone receptor beta: evidence for tissue-specific modulation of receptor function. *Embo j*. 1996; 15(12):3006–15. PMID: 8670802
65. Weiss RE, Forrest D, Pohlenz J, Cua K, Curran T, Refetoff S. Thyrotropin regulation by thyroid hormone in thyroid hormone receptor beta-deficient mice. *Endocrinology*. 1997; 138(9):3624–9. <https://doi.org/10.1210/endo.138.9.5412> PMID: 9275045
66. Gothe S, Wang Z, Ng L, Kindblom JM, Barros AC, Ohlsson C, et al. Mice devoid of all known thyroid hormone receptors are viable but exhibit disorders of the pituitary-thyroid axis, growth, and bone maturation. *Genes & development*. 1999; 13(10):1329–41.
67. Christoffolete MA, Ribeiro R, Singru P, Fekete C, da Silva WS, Gordon DF, et al. Atypical expression of type 2 iodothyronine deiodinase in thyrotrophs explains the thyroxine-mediated pituitary thyrotropin feedback mechanism. *Endocrinology*. 2006; 147(4):1735–43. <https://doi.org/10.1210/en.2005-1300> PMID: 16396983
68. Bianco AC, Salvatore D, Gereben B, Berry MJ, Larsen PR. Biochemistry, cellular and molecular biology, and physiological roles of the iodothyronine selenodeiodinases. *Endocrine reviews*. 2002; 23(1):38–89. <https://doi.org/10.1210/edrv.23.1.0455> PMID: 11844744
69. Gereben B, Zeold A, Dentice M, Salvatore D, Bianco AC. Activation and inactivation of thyroid hormone by deiodinases: local action with general consequences. *Cellular and molecular life sciences: CMLS*. 2008; 65(4):570–90. <https://doi.org/10.1007/s00018-007-7396-0> PMID: 17989921
70. Sugrue ML, Vella KR, Morales C, Lopez ME, Hollenberg AN. The thyrotropin-releasing hormone gene is regulated by thyroid hormone at the level of transcription in vivo. *Endocrinology*. 2010; 151(2):793–801. <https://doi.org/10.1210/en.2009-0976> PMID: 20032051

71. Fekete C, Lechan RM. Central regulation of hypothalamic-pituitary-thyroid axis under physiological and pathophysiological conditions. *Endocrine reviews*. 2014; 35(2):159–94. <https://doi.org/10.1210/er.2013-1087> PMID: 24423980
72. Nolan LA, Thomas CK, Levy A. Permissive effects of thyroid hormones on rat anterior pituitary mitotic activity. *The Journal of endocrinology*. 2004; 180(1):35–43. <https://doi.org/10.1677/joe.0.1800035> PMID: 14709142
73. Michaud JL, Rosenquist T, May NR, Fan CM. Development of neuroendocrine lineages requires the bHLH-PAS transcription factor SIM1. *Genes & development*. 1998; 12(20):3264–75.
74. Liu C, Goshu E, Wells A, Fan CM. Identification of the downstream targets of SIM1 and ARNT2, a pair of transcription factors essential for neuroendocrine cell differentiation. *The Journal of biological chemistry*. 2003; 278(45):44857–67. <https://doi.org/10.1074/jbc.M304489200> PMID: 12947113
75. Perello M, Friedman T, Paez-Espinosa V, Shen X, Stuart RC, Nillni EA. Thyroid hormones selectively regulate the posttranslational processing of prothyrotropin-releasing hormone in the paraventricular nucleus of the hypothalamus. *Endocrinology*. 2006; 147(6):2705–16. <https://doi.org/10.1210/en.2005-1609> PMID: 16497799
76. Lean AD, Ferland L, Drouin J, Kelly PA, Labrie F. Modulation of pituitary thyrotropin releasing hormone receptor levels by estrogens and thyroid hormones. *Endocrinology*. 1977; 100(6):1496–504. <https://doi.org/10.1210/endo-100-6-1496> PMID: 192537
77. Schomburg L, Bauer K. Thyroid hormones rapidly and stringently regulate the messenger RNA levels of the thyrotropin-releasing hormone (TRH) receptor and the TRH-degrading ectoenzyme. *Endocrinology*. 1995; 136(8):3480–5. <https://doi.org/10.1210/endo.136.8.7628384> PMID: 7628384
78. Krane IM, Spindel ER, Chin WW. Thyroid hormone decreases the stability and the poly(A) tract length of rat thyrotropin beta-subunit messenger RNA. *Molecular endocrinology (Baltimore, Md)*. 1991; 5(4):469–75.
79. Sanchez-Pacheco A, Palomino T, Aranda A. Negative regulation of expression of the pituitary-specific transcription factor GHF-1/Pit-1 by thyroid hormones through interference with promoter enhancer elements. *Molecular and cellular biology*. 1995; 15(11):6322–30. <https://doi.org/10.1128/mcb.15.11.6322> PMID: 7565785
80. Wood WM, Sarapura VD, Dowding JM, Woodmansee WW, Haakinson DJ, Gordon DF, et al. Early gene expression changes preceding thyroid hormone-induced involution of a thyrotrope tumor. *Endocrinology*. 2002; 143(2):347–59. <https://doi.org/10.1210/endo.143.2.8636> PMID: 11796486
81. Furumoto H, Ying H, Chandramouli GV, Zhao L, Walker RL, Meltzer PS, et al. An unliganded thyroid hormone beta receptor activates the cyclin D1/cyclin-dependent kinase/retinoblastoma/E2F pathway and induces pituitary tumorigenesis. *Molecular and cellular biology*. 2005; 25(1):124–35. <https://doi.org/10.1128/MCB.25.1.124-135.2005> PMID: 15601836
82. De Sousa SM, Earls P, McCormack AI. Pituitary hyperplasia: case series and literature review of an under-recognised and heterogeneous condition. *Endocrinology, diabetes & metabolism case reports*. 2015; 2015:150017.
83. Ingle E, Chappell D, Poon SY, Sarna MK, Beaumont JG, Williams JH, et al. Thyroid hormone receptor-interacting protein 1 modulates cytokine and nuclear hormone signaling in erythroid cells. *The Journal of biological chemistry*. 2001; 276(46):43428–34. <https://doi.org/10.1074/jbc.M106645200> PMID: 11544260
84. Park S, Han CR, Park JW, Zhao L, Zhu X, Willingham M, et al. Defective erythropoiesis caused by mutations of the thyroid hormone receptor alpha gene. *PLoS genetics*. 2017; 13(9):e1006991. <https://doi.org/10.1371/journal.pgen.1006991> PMID: 28910278
85. van Mullem AA, Visser TJ, Peeters RP. Clinical Consequences of Mutations in Thyroid Hormone Receptor-alpha1. *European thyroid journal*. 2014; 3(1):17–24. <https://doi.org/10.1159/000360637> PMID: 24847461
86. Kilby MD, Barber K, Hobbs E, Franklyn JA. Thyroid hormone action in the placenta. *Placenta*. 2005; 26(2–3):105–13. <https://doi.org/10.1016/j.placenta.2004.08.004> PMID: 15708111
87. Negro R, Schwartz A, Gismondi R, Tinelli A, Mangieri T, Stagnaro-Green A. Universal screening versus case finding for detection and treatment of thyroid hormonal dysfunction during pregnancy. *The Journal of clinical endocrinology and metabolism*. 2010; 95(4):1699–707. <https://doi.org/10.1210/jc.2009-2009> PMID: 20130074
88. De Groot L, Abalovich M, Alexander EK, Amino N, Barbour L, Cobin RH, et al. Management of thyroid dysfunction during pregnancy and postpartum: an Endocrine Society clinical practice guideline. *The Journal of clinical endocrinology and metabolism*. 2012; 97(8):2543–65. <https://doi.org/10.1210/jc.2011-2803> PMID: 22869843
89. Alexander EK, Pearce EN, Brent GA, Brown RS, Chen H, Dosiou C, et al. 2017 Guidelines of the American Thyroid Association for the Diagnosis and Management of Thyroid Disease During Pregnancy and the Postpartum. *Thyroid*. 2017; 27(3):315–89. <https://doi.org/10.1089/thy.2016.0457> PMID: 28056690